

SCORE Search Results Details for Application 10759514 and Search Result us-10-759-514-3.rng.

[Score Home](#)
[Page](#)

[Retrieve Application](#)
[List](#)

[SCORE System](#)
[Overview](#)

[SCORE](#)
[FAQ](#)

[Comments /](#)
[Suggestions](#)

This page gives you Search Results detail for the Application 10759514 and Search Result us-10-759-514-3.rng.

[start](#)

[Go Back to previous page](#)

GenCore version 5.1.8
Copyright (c) 1993 - 2006 Biocceleration Ltd.

OM nucleic - nucleic search, using sw model

Run on: May 7, 2006, 06:03:13 ; Search time 126.92 Seconds
(without alignments)
1260.261 Million cell updates/sec

Title: US-10-759-514-3
Perfect score: 24
Sequence: 1 ccgggagagccatagtgggtctgcg 24

Scoring table: IDENTITY_NUC
Gapop 10.0 , Gapext 1.0

Searched: 4996997 seqs, 3332346308 residues

Total number of hits satisfying chosen parameters: 9993994

Minimum DB seq length: 0
Maximum DB seq length: 2000000000

Post-processing: Minimum Match 0%
Maximum Match 100%
Listing first 45 summaries

Database : N_Geneseq_21:*
1: geneseqn1980s:*
2: geneseqn1990s:*
3: geneseqn2000s:*
4: geneseqn2001as:*
5: geneseqn2001bs:*
6: geneseqn2002as:*
7: geneseqn2002bs:*
8: geneseqn2003as:*
9: geneseqn2003bs:*
10: geneseqn2003cs:*
11: geneseqn2003ds:*
12: geneseqn2004as:*
13: geneseqn2004bs:*
14: geneseqn2005s:*

Pred. No. is the number of results predicted by chance to have a score greater than or equal to the score of the result being printed, and is derived by analysis of the total score distribution.

SUMMARIES

%

Result No.	Score	Query Match	Length	DB	ID	Description
1	24	100.0	24	10	ADD55637	Add55637 Oligonuc1
2	24	100.0	24	10	ADD55641	Add55641 Oligonuc1
3	24	100.0	24	14	ADV04767	Adv04767 Synthetic
4	24	100.0	24	14	ADV04754	Adv04754 Synthetic
5	24	100.0	24	14	ADZ75974	Adz75974 Hepatitis
6	24	100.0	24	14	ADZ75978	Adz75978 Hepatitis
c 7	24	100.0	27	2	AAQ64955	Aaq64955 Antisense
c 8	24	100.0	30	2	AAQ64950	Aaq64950 Antisense
9	24	100.0	37	2	AAX37631	Aax37631 HCV detec
10	24	100.0	50	3	AAA52575	Aaa52575 HCV RNA p
c 11	24	100.0	53	2	AAQ98103	Aaq98103 Label ext
c 12	24	100.0	70	2	AAT11268	Aat11268 Hepatitis
c 13	24	100.0	75	14	AEB94147	Aeb94147 Nucleic a
14	24	100.0	86	12	ADJ53747	Adj53747 HBV speci
15	24	100.0	102	4	AAC92379	Aac92379 HCV-RNA W
16	24	100.0	120	2	AAT69054	Aat69054 Hepatitis
c 17	24	100.0	120	14	AEB94144	Aeb94144 Nucleic a
18	24	100.0	120	14	AEB94143	Aeb94143 Nucleic a
19	24	100.0	131	14	ADW15169	Adw15169 HCV H77C
20	24	100.0	131	14	ADW15171	Adw15171 HCV from
21	24	100.0	131	14	ADW15170	Adw15170 HCV from
22	24	100.0	131	14	ADW15174	Adw15174 HCV from
23	24	100.0	131	14	ADW15172	Adw15172 HCV from
24	24	100.0	131	14	ADW15173	Adw15173 HCV from
c 25	24	100.0	140	2	AAT11269	Aat11269 Hepatitis
c 26	24	100.0	155	3	AAZ57775	Aaz57775 Hepatitis
27	24	100.0	159	2	AAQ43062	Aaq43062 -255 to -
28	24	100.0	159	2	AAQ43069	Aaq43069 -255 to -
29	24	100.0	159	2	AAQ43066	Aaq43066 -255 to -
30	24	100.0	159	2	AAQ43071	Aaq43071 -255 to -
31	24	100.0	177	2	AAQ79456	Aaq79456 HCV isola
32	24	100.0	177	2	AAQ68067	Aaq68067 HCV isola
33	24	100.0	177	2	AAQ79452	Aaq79452 HCV isola
34	24	100.0	177	2	AAQ79459	Aaq79459 HCV isola
35	24	100.0	177	2	AAQ79450	Aaq79450 HCV isola
36	24	100.0	177	2	AAQ79451	Aaq79451 HCV isola
37	24	100.0	177	2	AAQ68068	Aaq68068 HCV isola
38	24	100.0	177	2	AAQ79454	Aaq79454 HCV isola
39	24	100.0	177	2	AAQ68069	Aaq68069 HCV isola
40	24	100.0	177	2	AAQ68070	Aaq68070 HCV isola
41	24	100.0	177	2	AAQ79457	Aaq79457 HCV isola
42	24	100.0	177	2	AAQ68063	Aaq68063 HCV isola
43	24	100.0	177	2	AAQ79460	Aaq79460 HCV isola
44	24	100.0	177	2	AAQ79449	Aaq79449 HCV isola
45	24	100.0	177	2	AAQ79455	Aaq79455 HCV isola

ALIGNMENTS

RESULT 1

ADD55637

ID ADD55637 standard; DNA; 24 BP.

XX

AC ADD55637;

XX

DT 15-JAN-2004 (first entry)

XX

DE Oligonucleotide probe, PR2 #1, used to detect a HCV nucleic acid.

XX

KW HCV; fluorescent dye; fluorescent molecular beacon pair; lambda phage;

KW lambda phage-HCV hybrid amplicon; detection; diagnosis; HCV infection;
KW hepatitis; cirrhosis; antiviral therapy; probe; ss.
XX
OS Hepatitis C virus.
XX
PN US2003104582-A1.
XX
PD 05-JUN-2003.
XX
PF 04-DEC-2001; 2001US-00011855.
XX
PR 04-DEC-2001; 2001US-00011855.
XX
PA (BAUM/) BAUMANN R.
PA (HAMD/) HAMDAN H.
PA (LEWI/) LEWINSKI M.
XX
PI Baumann R, Hamdan H, Lewinski M;
XX
DR WPI; 2003-801237/75.
XX
PT Detecting hepatitis C virus (HCV) nucleic acid in a sample comprises
PT reverse transcribing and amplifying HCV nucleic acids with primer pair,
PT hybridizing amplicons with a labeled probe, and detecting a signal.
XX
PS Claim 1; Page 10; 11pp; English.
XX
CC The invention discloses a method for detecting the presence or amount of
CC Hepatitis C virus (HCV) nucleic acids in a sample comprising reverse
CC transcribing and amplifying any HCV nucleic acid present, reacting the
CC amplified nucleic acids with a probe in the presence of an enzyme that
CC cleaves the probe if specifically hybridised to HCV nucleic acids, and
CC detecting a signal from the probe. The detectable label is a fluorescent
CC dye or a fluorescent molecular beacon pair. Lambda phage HCV nucleic acid
CC hybrids are introduced into the test sample, reverse transcribed and
CC amplified using the pair of oligonucleotide primers to produce lambda
CC phage-HCV hybrid amplicons. The hybrids are hybridised to a control
CC oligonucleotide sequence (ADD55640) which is conjugated to 6-
CC carboxyfluorescein (FAM) and 6-carboxytetramethylrhodamine (TAMRA). The
CC test sample is chosen from serum, blood, plasma, cerebral spinal fluid,
CC synovial fluid, and urine. The nucleic acids are purified from the sample
CC prior to the reverse transcription and amplification step. The lambda
CC phage-HCV ribonucleic acid hybrids may be introduced into the test sample
CC prior to isolating nucleic acids from the sample. The method is useful
CC for detecting the presence or amount of hepatitis C virus (HCV) nucleic
CC acids in a test sample, for diagnosing HCV infection, which can lead to
CC chronic hepatitis and cirrhosis, for identification of individuals with
CC high viral replication, for monitoring patients on therapy and for
CC predicting whether antiviral therapy will be successful. The method is
CC specific and sensitive and exhibits a broad dynamic range of detection of
CC HCV nucleic acids and provides quantitative as well as qualitative
CC results. The sequence presented is an oligonucleotide probe used to
CC detect HCV nucleic acid.
XX
SQ Sequence 24 BP; 4 A; 6 C; 10 G; 4 T; 0 U; 0 Other;

Query Match 100.0%; Score 24; DB 10; Length 24;
Best Local Similarity 100.0%; Pred. No. 0.059;
Matches 24; Conservative 0; Mismatches 0; Indels 0; Gaps 0;

Qy 1 CCGGGAGAGCCATAGTGGTCTGCG 24
|||||
Db 1 CCGGGAGAGCCATAGTGGTCTGCG 24

RESULT 2

ADD55641

ID ADD55641 standard; DNA; 24 BP.

XX

AC ADD55641;

XX

DT 15-JAN-2004 (first entry)

XX

DE Oligonucleotide probe, PR2 #2, used to detect a HCV nucleic acid.

XX

KW HCV; fluorescent dye; fluorescent molecular beacon pair; lambda phage;

KW lambda phage-HCV hybrid amplicon; detection; diagnosis; HCV infection;

KW hepatitis; cirrhosis; antiviral therapy; probe; ss.

XX

OS Synthetic.

OS Hepatitis C virus.

XX

FH Key Location/Qualifiers

FT modified_base 1

FT /*tag= a

FT /mod_base= OTHER

FT /note= "OTHER= conjugated to 2'-chloro-7'-phenyl-1,4-

FT dichloro-6-carboxyfluorescein (VIC)"

FT modified_base 24

FT /*tag= b

FT /mod_base= OTHER

FT /note= "OTHER= conjugated to 6-

FT carboxytetramethylrhodamine (TAMRA)"

XX

PN US2003104582-A1.

XX

PD 05-JUN-2003.

XX

PF 04-DEC-2001; 2001US-00011855.

XX

PR 04-DEC-2001; 2001US-00011855.

XX

PA (BAUM/) BAUMANN R.

PA (HAMD/) HAMDAN H.

PA (LEWI/) LEWINSKI M.

XX

PI Baumann R, Hamdan H, Lewinski M;

XX

DR WPI; 2003-801237/75.

XX

PT Detecting hepatitis C virus (HCV) nucleic acid in a sample comprises

PT reverse transcribing and amplifying HCV nucleic acids with primer pair,

PT hybridizing amplicons with a labeled probe, and detecting a signal.

XX

PS Claim 5; Page 10; 11pp; English.

XX

CC The invention discloses a method for detecting the presence or amount of

CC Hepatitis C virus (HCV) nucleic acids in a sample comprising reverse

CC transcribing and amplifying any HCV nucleic acid present, reacting the

CC amplified nucleic acids with a probe in the presence of an enzyme that

CC cleaves the probe if specifically hybridised to HCV nucleic acids, and

CC detecting a signal from the probe. The detectable label is a fluorescent

CC dye or a fluorescent molecular beacon pair. Lambda phage HCV nucleic acid

CC hybrids are introduced into the test sample, reverse transcribed and

CC amplified using the pair of oligonucleotide primers to produce lambda

CC phage-HCV hybrid amplicons. The hybrids are hybridised to a control

CC oligonucleotide sequence (ADD55640) which is conjugated to 6-

CC carboxyfluorescein (FAM) and 6-carboxytetramethylrhodamine (TAMRA). The

CC test sample is chosen from serum, blood, plasma, cerebral spinal fluid,
 CC synovial fluid, and urine. The nucleic acids are purified from the sample
 CC prior to the reverse transcription and amplification step. The lambda
 CC phage-HCV ribonucleic acid hybrids may be introduced into the test sample
 CC prior to isolating nucleic acids from the sample. The method is useful
 CC for detecting the presence or amount of hepatitis C virus (HCV) nucleic
 CC acids in a test sample, for diagnosing HCV infection, which can lead to
 CC chronic hepatitis and cirrhosis, for identification of individuals with
 CC high viral replication, for monitoring patients on therapy and for
 CC predicting whether antiviral therapy will be successful. The method is
 CC specific and sensitive and exhibits a broad dynamic range of detection of
 CC HCV nucleic acids and provides quantitative as well as qualitative
 CC results. The sequence presented is a labelled oligonucleotide probe used
 CC to detect HCV nucleic acid.

XX

SQ Sequence 24 BP; 4 A; 6 C; 10 G; 4 T; 0 U; 0 Other;

Query Match 100.0%; Score 24; DB 10; Length 24;
 Best Local Similarity 100.0%; Pred. No. 0.059;
 Matches 24; Conservative 0; Mismatches 0; Indels 0; Gaps 0;

Qy 1 CCGGGAGAGCCATAGTGGTCTGCG 24
 |||||
 Db 1 CCGGGAGAGCCATAGTGGTCTGCG 24

RESULT 3

ADV04767

ID ADV04767 standard; DNA; 24 BP.

XX

AC ADV04767;

XX

DT 24-FEB-2005 (first entry)

XX

DE Synthetic PCR primer #18.

XX

KW Virucide; hepatitis C virus infection; ss; replicon; PCR; primer.

XX

OS Synthetic.

XX

PN WO2004104198-A1.

XX

PD 02-DEC-2004.

XX

PF 25-NOV-2003; 2003WO-JP015038.

XX

PR 26-MAY-2003; 2003JP-00148242.

PR 19-SEP-2003; 2003JP-00329115.

XX

PA (TORA) TORAY IND INC.

PA (TOKM-) TOKYO METROPOLITAN ORG MEDICAL RES.

PA (UYMA-) UNIV MAINZ GUTENBERG JOHANNES.

XX

PI Wakita T, Kato T, Date T;

XX

DR WPI; 2005-013292/01.

XX

PT Novel replicon RNA, having sequence of 5' and 3' untranslated region and
 PT base sequence encoding NS3, NS4A, NS4B, NS5A and NS5B proteins on genomic
 PT RNA of hepatitis C virus of genotype 2a, useful for treating hepatitis C
 PT virus infection.

XX

PS Example 7; SEQ ID NO 33; 197pp; Japanese.

XX

CC The invention relates to replicon RNA from genotype 2a of hepatitis C
 CC virus comprising a 5' untranslated region, a base sequence encoding NS3
 CC protein, NS4A protein, NS4B protein, NS5A protein and NS5B protein, and a
 CC 3' untranslated region. The invention also relates to a cell capable of
 CC reproducing the replicon involving transducing the replicon RNA to a
 CC cell, a method of producing a hepatitis C virus protein, a method of
 CC screening a substance that promotes or suppresses the reproduction of
 CC hepatitis C virus, involving culturing the replicon reproducing cell in
 CC the presence of a test substance, and detecting the reproduction of
 CC replicon RNA in the culture. Virucide. The replicon RNA is useful for
 CC producing a replicon reproduction cell and for increasing the
 CC reproduction efficiency of replicon RNA of hepatitis C virus of genotype
 CC 2a. The cell and the replicon RNA are useful for producing a therapeutic
 CC agent or a diagnostic agent for hepatitis C virus infection, for
 CC producing a vaccine against hepatitis C virus infection and for screening
 CC a substance that promotes or suppresses the reproduction of hepatitis C
 CC virus. This sequence represents a PCR primer used in the scope of the
 CC invention.

XX

SQ Sequence 24 BP; 4 A; 6 C; 10 G; 4 T; 0 U; 0 Other;

Query Match 100.0%; Score 24; DB 14; Length 24;
 Best Local Similarity 100.0%; Pred. No. 0.059;
 Matches 24; Conservative 0; Mismatches 0; Indels 0; Gaps 0;

Qy 1 CCGGGAGAGCCATAGTGGTCTGCG 24
 |||||
 Db 1 CCGGGAGAGCCATAGTGGTCTGCG 24

RESULT 4

ADV04754

ID ADV04754 standard; DNA; 24 BP.

XX

AC ADV04754;

XX

DT 24-FEB-2005 (first entry)

XX

DE Synthetic PCR primer #5.

XX

KW Virucide; hepatitis C virus infection; ss; replicon; PCR; primer.

XX

OS Synthetic.

XX

PN WO2004104198-A1.

XX

PD 02-DEC-2004.

XX

PF 25-NOV-2003; 2003WO-JP015038.

XX

PR 26-MAY-2003; 2003JP-00148242.

XX

PR 19-SEP-2003; 2003JP-00329115.

XX

PA (TORA) TORAY IND INC.

PA (TOKM-) TOKYO METROPOLITAN ORG MEDICAL RES.

PA (UYMA-) UNIV MAINZ GUTENBERG JOHANNES.

XX

PI Wakita T, Kato T, Date T;

XX

DR WPI; 2005-013292/01.

XX

PT Novel replicon RNA, having sequence of 5' and 3' untranslated region and
 PT base sequence encoding NS3, NS4A, NS4B, NS5A and NS5B proteins on genomic
 PT RNA of hepatitis C virus of genotype 2a, useful for treating hepatitis C

PT virus infection.
 XX
 PS Example 5; SEQ ID NO 20; 197pp; Japanese.
 XX
 CC The invention relates to replicon RNA from genotype 2a of hepatitis C
 CC virus comprising a 5' untranslated region, a base sequence encoding NS3
 CC protein, NS4A protein, NS4B protein, NS5A protein and NS5B protein, and a
 CC 3' untranslated region. The invention also relates to a cell capable of
 CC reproducing the replicon involving transducing the replicon RNA to a
 CC cell, a method of producing a hepatitis C virus protein, a method of
 CC screening a substance that promotes or suppresses the reproduction of
 CC hepatitis C virus, involving culturing the replicon reproducing cell in
 CC the presence of a test substance, and detecting the reproduction of
 CC replicon RNA in the culture. Virucide. The replicon RNA is useful for
 CC producing a replicon reproduction cell and for increasing the
 CC reproduction efficiency of replicon RNA of hepatitis C virus of genotype
 CC 2a. The cell and the replicon RNA are useful for producing a therapeutic
 CC agent or a diagnostic agent for hepatitis C virus infection, for
 CC producing a vaccine against hepatitis C virus infection and for screening
 CC a substance that promotes or suppresses the reproduction of hepatitis C
 CC virus. This sequence represents a PCR primer used in the scope of the
 CC invention.
 XX
 SQ Sequence 24 BP; 4 A; 6 C; 10 G; 4 T; 0 U; 0 Other;

Query Match 100.0%; Score 24; DB 14; Length 24;
 Best Local Similarity 100.0%; Pred. No. 0.059;
 Matches 24; Conservative 0; Mismatches 0; Indels 0; Gaps 0;

Qy 1 CCGGGAGAGCCATAGTGGTCTGCG 24
 |||||
 Db 1 CCGGGAGAGCCATAGTGGTCTGCG 24

RESULT 5

ADZ75974

ID ADZ75974 standard; DNA; 24 BP.

XX

AC ADZ75974;

XX

DT 14-JUL-2005 (first entry)

XX

DE Hepatitis C virus specific probe SEQ ID NO:3.

XX

KW DNA detection; RNA detection; hepatitis C virus infection;

KW antiinflammatory; hepatotropic; virucide; probe; ss.

XX

OS Hepatitis C virus.

OS Synthetic.

XX

PN US2005100889-A1.

XX

PD 12-MAY-2005.

XX

PF 13-OCT-2004; 2004US-00964302.

XX

PR 04-DEC-2001; 2001US-00011855.

XX

PA (QUES-) QUEST DIAGNOSTICS INVESTMENTS INC.

XX

PI Baumann R, Hamdan H, Lewinski M;

XX

DR WPI; 2005-345387/35.

XX

PT Detecting Hepatitis C virus (HCV) nucleic acids in a test sample, for
PT diagnosing HCV infection, comprises using oligonucleotide primers and
PT probes to amplify HCV and/or control nucleic acid sequences present in
PT the sample.
XX
PS Claim 1; SEQ ID NO 3; 13pp; English.
XX
CC The invention relates to a method for detecting the presence or amount of
CC Hepatitis C virus (HCV) nucleic acids in a test sample. The method
CC comprises: (a) introducing lambda phage-HCV nucleic acid hybrids into the
CC test sample; (b) reverse transcribing and amplifying: (i) HCV nucleic
CC acid if present in the sample and using a pair of oligonucleotide primers
CC having the sequences set forth in ADZ75972 and ADZ75973, to generate HCV
CC amplicons; and (ii) lambda phage-HCV hybrid nucleic acid using a pair of
CC oligonucleotide primers having the sequences set forth in ADZ75972 and
CC ADZ75973, to generate a lambda phage-HCV hybrid amplicons; (c)
CC hybridizing the HCV amplicons with an oligonucleotide probe comprising
CC the 24 bp sequence of ADZ75974 in the presence of an enzyme that cleaves
CC the probe when the probe hybridizes to the HCV nucleic acids, where the
CC probe is conjugated to a first detectable label that generates a
CC detectable signal upon the cleavage; (d) hybridizing the lambda phage-HCV
CC hybrid amplicons to a control oligonucleotide probe in the presence of an
CC enzyme that cleaves the control oligonucleotide probe when the control
CC probe hybridizes to the lambda phage-HCV hybrid amplicons, where the
CC control probe is conjugated to a second detectable label that generates a
CC detectable signal upon the cleavage; and (e) detecting a signal from the
CC first and second detectable labels, where the signal from the first
CC detectable label indicates the presence or amount of HCV nucleic acids in
CC the test sample. Also described is a composition of one or more
CC substantially purified oligonucleotides comprising the nucleotide
CC sequences mentioned in the specification, and a kit comprising materials
CC for performing the new method. The method and composition are useful for
CC diagnosing HCV infection or for qualitatively and quantitatively
CC detecting hepatitis C viral nucleic acids in a test sample. The present
CC sequence represents an HCV specific probe which is used in an example
CC from the present invention.
XX
SQ Sequence 24 BP; 4 A; 6 C; 10 G; 4 T; 0 U; 0 Other;

Query Match 100.0%; Score 24; DB 14; Length 24;
Best Local Similarity 100.0%; Pred. No. 0.059;
Matches 24; Conservative 0; Mismatches 0; Indels 0; Gaps 0;

Qy 1 CCGGGAGAGCCATAGTGGTCTGCG 24
|||||
Db 1 CCGGGAGAGCCATAGTGGTCTGCG 24

RESULT 6
ADZ75978
ID ADZ75978 standard; DNA; 24 BP.
XX
AC ADZ75978;
XX
DT 14-JUL-2005 (first entry)
XX
DE Hepatitis C virus specific probe SEQ ID NO:7.
XX
KW DNA detection; RNA detection; hepatitis C virus infection;
KW antiinflammatory; hepatotropic; virucide; probe; ss.
XX
OS Hepatitis C virus.
OS Synthetic.
XX

```

FH   Key                Location/Qualifiers
FT   misc_binding       1
FT                               /*tag=  a
FT                               /bound_moiety= "2'-chloro-7'-phenyl-1-4-dichloro-6-
FT                               carboxyfluorescein (VIC)"
FT   misc_binding       24
FT                               /*tag=  b
FT                               /bound_moiety= "6-carboxytetramethylrhodamine (TAMRA)"
XX
PN   US2005100889-A1.
XX
PD   12-MAY-2005.
XX
PF   13-OCT-2004; 2004US-00964302.
XX
PR   04-DEC-2001; 2001US-00011855.
XX
PA   (QUES-) QUEST DIAGNOSTICS INVESTMENTS INC.
XX
PI   Baumann R,  Hamdan H,  Lewinski M;
XX
DR   WPI; 2005-345387/35.
XX
PT   Detecting Hepatitis C virus (HCV) nucleic acids in a test sample, for
PT   diagnosing HCV infection, comprises using oligonucleotide primers and
PT   probes to amplify HCV and/or control nucleic acid sequences present in
PT   the sample.
XX
PS   Example 2; SEQ ID NO 7; 13pp; English.
XX
CC   The invention relates to a method for detecting the presence or amount of
CC   Hepatitis C virus (HCV) nucleic acids in a test sample. The method
CC   comprises: (a) introducing lambda phage-HCV nucleic acid hybrids into the
CC   test sample; (b) reverse transcribing and amplifying: (i) HCV nucleic
CC   acid if present in the sample and using a pair of oligonucleotide primers
CC   having the sequences set forth in ADZ75972 and ADZ75973, to generate HCV
CC   amplicons; and (ii) lambda phage-HCV hybrid nucleic acid using a pair of
CC   oligonucleotide primers having the sequences set forth in ADZ75972 and
CC   ADZ75973, to generate a lambda phage-HCV hybrid amplicons; (c)
CC   hybridizing the HCV amplicons with an oligonucleotide probe comprising
CC   the 24 bp sequence of ADZ75974 in the presence of an enzyme that cleaves
CC   the probe when the probe hybridizes to the HCV nucleic acids, where the
CC   probe is conjugated to a first detectable label that generates a
CC   detectable signal upon the cleavage; (d) hybridizing the lambda phage-HCV
CC   hybrid amplicons to a control oligonucleotide probe in the presence of an
CC   enzyme that cleaves the control oligonucleotide probe when the control
CC   probe hybridizes to the lambda phage-HCV hybrid amplicons, where the
CC   control probe is conjugated to a second detectable label that generates a
CC   detectable signal upon the cleavage; and (e) detecting a signal from the
CC   first and second detectable labels, where the signal from the first
CC   detectable label indicates the presence or amount of HCV nucleic acids in
CC   the test sample. Also described is a composition of one or more
CC   substantially purified oligonucleotides comprising the nucleotide
CC   sequences mentioned in the specification, and a kit comprising materials
CC   for performing the new method. The method and composition are useful for
CC   diagnosing HCV infection or for qualitatively and quantitatively
CC   detecting hepatitis C viral nucleic acids in a test sample. The present
CC   sequence represents an HCV specific probe which is used in an example
CC   from the present invention.
XX
SQ   Sequence 24 BP; 4 A; 6 C; 10 G; 4 T; 0 U; 0 Other;

```

```

Query Match          100.0%;  Score 24;  DB 14;  Length 24;
Best Local Similarity 100.0%;  Pred. No. 0.059;

```

Matches 24; Conservative 0; Mismatches 0; Indels 0; Gaps 0;

```
Qy      1 CCGGGAGAGCCATAGTGGTCTGCG 24
          |||
Db      1 CCGGGAGAGCCATAGTGGTCTGCG 24
```

RESULT 7

AAQ64955/c

ID AAQ64955 standard; DNA; 27 BP.

XX

AC AAQ64955;

XX

DT 19-DEC-1994 (first entry)

XX

DE Antisense oligonucleotide complementary to Hepatitis C Virus genome.

XX

KW Hepatitis C Virus; Non-A, non-B hepatitis virus; HCV; antisense; therapy;
inhibition; viral protein precursor; ss.

XX

OS Synthetic.

XX

PN CA2104649-A.

XX

PD 26-FEB-1994.

XX

PF 23-AUG-1993; 93CA-02104649.

XX

PR 25-AUG-1992; 92JP-00248796.

PR 03-MAR-1993; 93JP-00042736.

XX

PA (SEKI/) SEKI M.

XX

PI Seki M, Honda Y, Yamada E;

XX

DR WPI; 1994-151836/19.

XX

PT Anti:sense oligo:nucleotide(s) complementary to the hepatitis C virus
genome - are useful as antiviral agents.

XX

PS Claim 5; Page 82; 262pp; English.

XX

CC This oligonucleotide is an example of a preferred antisense compound i.e.
CC it has a base sequence of 15-30 bases which is included within the 31
CC bases from C at position 150 to G at position 180 of AAQ64913 and which
CC contains at least 6 bases from C at position 175 to G at position 180.
CC The antisense oligonucleotide is useful for inhibiting translation of HCV
CC genes

XX

SQ Sequence 27 BP; 5 A; 11 C; 7 G; 4 T; 0 U; 0 Other;

Query Match 100.0%; Score 24; DB 2; Length 27;

Best Local Similarity 100.0%; Pred. No. 0.06;

Matches 24; Conservative 0; Mismatches 0; Indels 0; Gaps 0;

```
Qy      1 CCGGGAGAGCCATAGTGGTCTGCG 24
          |||
Db     25 CCGGGAGAGCCATAGTGGTCTGCG 2
```

RESULT 8

AAQ64950/c

ID AAQ64950 standard; DNA; 30 BP.

XX

AC AAQ64950;
 XX
 DT 19-DEC-1994 (first entry)
 XX
 DE Antisense oligonucleotide complementary to Hepatitis C Virus genome.
 XX
 KW Hepatitis C Virus; Non-A, non-B hepatitis virus; HCV; antisense; therapy;
 KW inhibition; viral protein precursor; ss.
 XX
 OS Synthetic.
 XX
 PN CA2104649-A.
 XX
 PD 26-FEB-1994.
 XX
 PF 23-AUG-1993; 93CA-02104649.
 XX
 PR 25-AUG-1992; 92JP-00248796.
 PR 03-MAR-1993; 93JP-00042736.
 XX
 PA (SEKI/) SEKI M.
 XX
 PI Seki M, Honda Y, Yamada E;
 XX
 DR WPI; 1994-151836/19.
 XX
 PT Anti:sense oligo:nucleotide(s) complementary to the hepatitis C virus
 PT genome - are useful as antiviral agents.
 XX
 PS Claim 5; Page 80; 262pp; English.
 XX
 CC This oligonucleotide is an example of a preferred antisense compound i.e.
 CC it has a base sequence of 15-30 bases which is included within the 31
 CC bases from C at position 150 to G at position 180 of AAQ64913 and which
 CC contains at least 6 bases from C at position 151 to C at position 156.
 CC The antisense oligonucleotide is useful for inhibiting translation of HCV
 CC genes
 XX
 SQ Sequence 30 BP; 5 A; 11 C; 10 G; 4 T; 0 U; 0 Other;

Query Match 100.0%; Score 24; DB 2; Length 30;
 Best Local Similarity 100.0%; Pred. No. 0.061;
 Matches 24; Conservative 0; Mismatches 0; Indels 0; Gaps 0;

Qy 1 CCGGGAGAGCCATAGTGGTCTGCG 24
 |||||
 Db 25 CCGGGAGAGCCATAGTGGTCTGCG 2

RESULT 9

AAX37631

ID AAX37631 standard; DNA; 37 BP.

XX

AC AAX37631;

XX

DT 08-JUL-1999 (first entry)

XX

DE HCV detecting primer #1.

XX

KW Detection; HCV; real time; PCR; reporter; fluorescent; primer; quencher;

KW fluorescence resonance energy transfer; ss;

XX

OS Synthetic.

OS Hepatitis C virus; Virus.

XX
 PN JP11103899-A.
 XX
 PD 20-APR-1999.
 XX
 PF 30-SEP-1997; 97JP-00283042.
 XX
 PR 30-SEP-1997; 97JP-00283042.
 XX
 PA (TOKR-) ZH TOKYOTO RINSHO IGAKU SOGO KENKYUSHO.
 PA (SRLS-) SRL KK.
 XX
 DR WPI; 1999-305862/26.
 XX
 PT Measurement of HCV gene using real time detecting PCR and primer and
 PT probe - is highly sensitive.
 XX
 PS Claim 1; Page 6; 8pp; Japanese.
 XX
 CC This invention describes a method for the measurement of an HCV gene by a
 CC real time detecting PCR. The invention also describes a method where a
 CC reporter fluorescent colour and a quencher fluorescent colour are
 CC combined to an oligonucleotide and the fluorescence of the reporter
 CC fluorescent colour is controlled by fluorescence resonance energy
 CC transfer. The method can measure HCV exactly with high sensitivity
 XX
 SQ Sequence 37 BP; 6 A; 15 C; 11 G; 5 T; 0 U; 0 Other;

Query Match 100.0%; Score 24; DB 2; Length 37;
 Best Local Similarity 100.0%; Pred. No. 0.063;
 Matches 24; Conservative 0; Mismatches 0; Indels 0; Gaps 0;

Qy 1 CCGGGAGAGCCATAGTGGTCTGCG 24
 ||||||||||||||||||
 Db 10 CCGGGAGAGCCATAGTGGTCTGCG 33

RESULT 10

AAA52575

ID AAA52575 standard; DNA; 50 BP.

XX

AC AAA52575;

XX

DT 27-SEP-2000 (first entry)

XX

DE HCV RNA promoter primer, SEQ ID NO:13.

XX

KW Oligonucleotide; HCV genomic RNA; detection; amplification;

KW reverse transcription inhibition; translation inhibition; antiviral;

KW gene therapy; sense; promoter primer; reverse transcription-PCR;

KW RT-PCR primer; ss.

XX

OS Hepatitis C virus.

XX

PN EP1002878-A2.

XX

PD 24-MAY-2000.

XX

PF 18-NOV-1999; 99EP-00122092.

XX

PR 19-NOV-1998; 98JP-00329874.

XX

PA (TOYJ) TOSOH CORP.

XX

PI Toshiki T, Takahiko I, Juichi S;
 XX
 DR WPI; 2000-352431/31.
 XX
 PT Hepatitis C virus RNA-binding single-stranded oligo DNAs useful as
 PT reagents for gene diagnosis involving cleavage, amplification and
 PT detection of RNA and as an inhibitory drugs.
 XX
 PS Example 5; Page 15; 21pp; English.
 XX
 CC The invention relates to single-stranded antisense oligodeoxynucleotides
 CC (AAA52563-A52568) which bind to various sites on the hepatitis C virus
 CC (HCV) RNA genome, and to sense oligodeoxynucleotides (AAA52569-A52571)
 CC corresponding to sites on the HCV genome. The oligonucleotides are useful
 CC as primers in RT-PCR (reverse transcription-PCR) and the sense
 CC oligonucleotides may also be used as promoter primers. The antisense
 CC oligonucleotides may be used to inhibit translation or reverse
 CC transcription of HCV RNA and may be used as probes for detection of HCV
 CC RNA. Additionally, the antisense oligos may be linked to an RNA-cleaving
 CC moiety to target single-stranded RNA cleavage or RNA heteroduplex
 CC cleavage. The invention also encompasses methods of identifying and
 CC preparing single-stranded oligodeoxynucleotides which bind to target
 CC RNAs. The single-stranded oligodeoxynucleotides are useful as reagents
 CC for genetic diagnosis involving cleavage, amplification and detection of
 CC HCV RNA (as primers and probes), and as inhibitors of reverse
 CC transcription or translation of HCV RNA. Sequences AAA42573-A52576
 CC represent HCV RNA promoter primers used in an exemplification of the
 CC invention
 XX
 SQ Sequence 50 BP; 15 A; 9 C; 15 G; 11 T; 0 U; 0 Other;

Query Match 100.0%; Score 24; DB 3; Length 50;
 Best Local Similarity 100.0%; Pred. No. 0.066;
 Matches 24; Conservative 0; Mismatches 0; Indels 0; Gaps 0;

Qy 1 CCGGGAGAGCCATAGTGGTCTGCG 24
 ||||||||||||||||||
 Db 26 CCGGGAGAGCCATAGTGGTCTGCG 49

RESULT 11

AAQ98103/c

ID AAQ98103 standard; DNA; 53 BP.

XX

AC AAQ98103;

XX

DT 05-FEB-1996 (first entry)

XX

DE Label extender probe used in an improved sandwich hybridisation assay.

XX

KW Probe; nucleotide; solution phase sandwich hybridisation assay;
 KW competitive; analyte binding sequence; background signal reduction; ss.

XX

OS Synthetic.

XX

PN WO9516055-A1.

XX

PD 15-JUN-1995.

XX

PF 07-DEC-1994; 94WO-US014119.

XX

PR 08-DEC-1993; 93US-00164388.

XX

PA (CHIR) CHIRON CORP.

XX
 PI Urdea MS, Fultz T, Warner BD, Collins M;
 XX
 DR WPI; 1995-224335/29.
 XX
 PT Soln. phase sandwich hybridisation assays for nucleic acid(s) - with
 PT capture extender molecules or competitive oligo:nucleotide(s) to minimise
 PT background signal, increasing sensitivity and selectivity.
 XX
 PS Example 1; Page 33; 86pp; English.
 XX
 CC AAQ98100-Q98105 are label extender probes (LEs) used in a variation of a
 CC new improved method of a solution phase sandwich hybridisation assay in
 CC which LEs are used with a capture probe (CP). One label extender probe
 CC binds the target DNA and another binds to a labelled probe (LP). The new
 CC method minimises background signals (caused by non-specific
 CC hybridisation), this improves both sensitivity and selectivity of the
 CC assay without increasing cost or time
 XX
 SQ Sequence 53 BP; 9 A; 15 C; 20 G; 9 T; 0 U; 0 Other;

Query Match 100.0%; Score 24; DB 2; Length 53;
 Best Local Similarity 100.0%; Pred. No. 0.066;
 Matches 24; Conservative 0; Mismatches 0; Indels 0; Gaps 0;

QY 1 CCGGGAGAGCCATAGTGGTCTGCG 24
 ||||||||||||||||||
 Db 44 CCGGGAGAGCCATAGTGGTCTGCG 21

RESULT 12

AAT11268/c

ID AAT11268 standard; RNA; 70 BP.

XX

AC AAT11268;

XX

DT 26-JUN-1996 (first entry)

XX

DE Hepatitis C virus partial 5'-UTR antisense RNA AS3.

XX

KW Antisense; therapy; complementary; HCV; 5'-untranslated region;
 KW hepatitis C virus; inhibition; infection; treatment; stem-loop;
 KW clone 2-1; ss.

XX

OS Hepatitis C virus.

XX

PN JP07303485-A.

XX

PD 21-NOV-1995.

XX

PF 13-MAY-1994; 94JP-00124609.

XX

PR 13-MAY-1994; 94JP-00124609.

XX

PA (TOFU) TONEN CORP.

XX

DR WPI; 1996-035187/04.

XX

PT Hepatitis C virus (HCV) anti:sense RNA - inhibits HCV structural gene
 PT expression in vivo for treatment of HCV infection.

XX

PS Claim 2; Page 9; 12pp; Japanese.

XX

CC The present sequence is a specifically claimed example of RNA that is

CC complementary (i.e. antisense) to part of the 5'-untranslated region of
CC the hepatitis C virus genome sequence contained in clone 2-1. The 5'-UTR
CC includes several stem-loop sequences. The antisense RNA is useful for
CC inhibiting expression of HCV structural genes and thereby inhibiting
CC viral replication in vivo. The antisense therapy can be used in addition
CC to conventional interferon treatment of HCV infections

XX

SQ Sequence 70 BP; 10 A; 21 C; 25 G; 0 T; 14 U; 0 Other;

Query Match 100.0%; Score 24; DB 2; Length 70;
Best Local Similarity 100.0%; Pred. No. 0.069;
Matches 24; Conservative 0; Mismatches 0; Indels 0; Gaps 0;

Qy 1 CCGGGAGAGCCATAGTGGTCTGCG 24
|||||||
Db 51 CCGGGAGAGCCATAGTGGTCTGCG 28

RESULT 13

AEB94147/c

ID AEB94147 standard; DNA; 75 BP.

XX

AC AEB94147;

XX

DT 06-OCT-2005 (first entry)

XX

DE Nucleic acid detection method target sequence LTar-HCV.

XX

KW ss; probe; microorganism detection; DNA detection; RNA detection.

XX

OS Hepatitis C virus.

XX

PN WO2005071401-A2.

XX

PD 04-AUG-2005.

XX

PF 14-JAN-2005; 2005WO-US001378.

XX

PR 15-JAN-2004; 2004US-0536978P.

XX

PA (CHIR) CHIRON CORP.

XX

PI Shyamala V, Nguyen SH;

XX

DR WPI; 2005-564246/57.

XX

PT Detecting a nucleic acid target sequence comprises contacting the sample
PT with a capture probe conjugate, a reporter probe and a detectable
PT reporter moiety.

XX

PS Example 1; SEQ ID NO 22; 42pp; English.

XX

CC The invention relates to a method of detecting a nucleic acid target
CC sequence in a sample comprising contacting the sample with a capture
CC probe conjugate, a reporter probe and a detectable reporter moiety. The
CC method comprises: (i) contacting the sample with a capture probe
CC conjugate, a reporter probe and a detectable reporter moiety, under
CC conditions allowing formation of a complex including the nucleic acid
CC target sequence, if present, the capture probe conjugate, the reporter
CC probe and the detectable reporter moiety, where the capture probe
CC conjugate and the reporter probe each comprise an oligonucleotide that is
CC capable of specifically hybridizing to the nucleic acid target sequence
CC and where the capture probe conjugate and the reporter probe do not
CC hybridize to the same or overlapping regions of the nucleic acid target

CC sequence. The capture probe conjugate comprises a substrate having a
 CC distinguishable spectral signal signature that uniquely identifies the
 CC capture probe conjugate, where the reporter probe comprises one member of
 CC a binding pair and the detectable reporter moiety comprises the other
 CC member of the binding pair, and where the detectable reporter moiety
 CC comprises a detectable label; and (ii) detecting a signal from the
 CC capture probe conjugate substrate and the detectable reporter moiety
 CC label individually from each complex so formed. Also included is a kit
 CC for a multiplex assay for detecting the presence of nucleic acid target
 CC sequences in a sample, comprising: capture probe conjugates, reporter
 CC probes and a detectable reporter moiety, where each member of the capture
 CC probe conjugates is specific for one member of the target sequences (the
 CC specific capture probe conjugate) and each member of the reporter probes
 CC is specific for one member of the target sequences (the specific reporter
 CC probe). The nucleic acid target sequence is a sequence from a pathogen
 CC nucleic acid, where the pathogen is a virus selected from HIV, HBV, HCV,
 CC HAV, parvovirus B19, West Nile Virus, hantavirus or SARS. The method and
 CC kit are useful for detecting a nucleic acid target sequence in a sample.
 CC The present sequence represents a nucleic acid detection method target
 CC sequence.

XX

SQ Sequence 75 BP; 14 A; 24 C; 23 G; 14 T; 0 U; 0 Other;

Query Match 100.0%; Score 24; DB 14; Length 75;
 Best Local Similarity 100.0%; Pred. No. 0.07;
 Matches 24; Conservative 0; Mismatches 0; Indels 0; Gaps 0;

Qy 1 CCGGGAGAGCCATAGTGGTCTGCG 24
 |||||
 Db 27 CCGGGAGAGCCATAGTGGTCTGCG 4

RESULT 14

ADJ53747

ID ADJ53747 standard; DNA; 86 BP.

XX

AC ADJ53747;

XX

DT 06-MAY-2004 (first entry)

XX

DE HBV specific molecular beacon target #16.

XX

KW ss; capture oligonucleotide; HBV; HIV-1; HCV; donated blood screening.

XX

OS Hepatitis B virus.

XX

PN WO2003106714-A1.

XX

PD 24-DEC-2003.

XX

PF 13-JUN-2003; 2003WO-US018993.

XX

PR 14-JUN-2002; 2002US-0389393P.

XX

PA (GENP-) GEN-PROBE INC.

XX

PI Linnen JM, Kolk DP, Dockter JM, Getman DK, Yoshimura T;

PI Ho-Sing-Loy M, Stringfellow LA;

XX

DR WPI; 2004-082210/08.

XX

PT Capture oligonucleotide composition useful for detection of hepatitis B
 PT virus (HBV), comprising polynucleotide having HBV-complementary sequence
 PT which is immobilized on solid support.

XX
 PS Example 10; SEQ ID NO 141; 112pp; English.
 XX
 CC The invention relates to a capture oligonucleotide composition comprising
 CC an hepatitis B virus (HBV)-complementary sequence polynucleotide
 CC immobilised to a solid support. The composition is useful for detecting
 CC nucleic acids of HBV and/of HIV-1 and/or HCV in biological sample such as
 CC blood, serum, plasma or other body fluid or tissue to be tested. The
 CC composition can be used either in diagnostic application or for screening
 CC donated blood and that products or other tissues that may contain
 CC infectious particles. The composition facilitates detection of very low
 CC levels of HBV nucleic acids. The composition allows selective detection
 CC of nucleic acids of HBV and/or HIV and/or HCV. The present sequence is
 CC used in the exemplification of the invention.
 XX
 SQ Sequence 86 BP; 17 A; 26 C; 27 G; 16 T; 0 U; 0 Other;

Query Match 100.0%; Score 24; DB 12; Length 86;
 Best Local Similarity 100.0%; Pred. No. 0.071;
 Matches 24; Conservative 0; Mismatches 0; Indels 0; Gaps 0;

Qy 1 CCGGGAGAGCCATAGTGGTCTGCG 24
 |||||
 Db 38 CCGGGAGAGCCATAGTGGTCTGCG 61

RESULT 15

AAC92379

ID AAC92379 standard; RNA; 102 BP.

XX

AC AAC92379;

XX

DT 26-MAR-2001 (first entry)

XX

DE HCV-RNA WQ-RNA nucleotide sequence SEQ ID NO:7.

XX

KW Amplification; analysis; RNA transcription; RNA polymerase; detection;

KW bacterium; virus; foodstuff; soil; environmental water; seawater;

KW house dust; ss.

XX

OS Synthetic.

XX

PN WO200075371-A1.

XX

PD 14-DEC-2000.

XX

PF 05-JUN-2000; 2000WO-JP003647.

XX

PR 04-JUN-1999; 99JP-00157653.

XX

PA (TOYJ) TOSOH CORP.

XX

PI Ishizuka T, Ishiguro T, Saitoh J, Sakai T;

XX

DR WPI; 2001-061738/07.

XX

PT Potentiated method for amplification of nucleic acids for detection in

PT biological samples.

XX

PS Example 3; Page 38; 42pp; Japanese.

XX

CC The present invention describes a method for amplifying a specific RNA

CC from a sample. The method comprises generating from the template RNA in

CC the sample a two-stranded DNA, which contains sequences complementary to

CC and homologous with the RNA sequence, and a transcription promoter. RNA
CC is transcribed from the two-stranded DNA using an RNA polymerase, in as
CC reaction mixture comprising inosine triphosphate (ITP), Adenosine
CC triphosphate (ATP), uridine triphosphate (UTP), cytidine triphosphate
CC (CTP) and guanosine triphosphate (GTP), and the reaction cycle is
CC repeated as necessary. The method can be used for the detection and assay
CC of specific RNA sequences, such as those occurring in bacteria and
CC viruses, in samples such as foodstuffs, soil, environmental waters,
CC seawater and house dust. The addition of ITP increases the efficiency of
CC the amplification reaction. The present sequence represents an
CC oligonucleotide used in an example from the present invention, for the
CC exemplification of the method of the invention

XX

SQ Sequence 102 BP; 23 A; 29 C; 25 G; 0 T; 25 U; 0 Other;

Query Match 100.0%; Score 24; DB 4; Length 102;
Best Local Similarity 83.3%; Pred. No. 0.073;
Matches 20; Conservative 4; Mismatches 0; Indels 0; Gaps 0;

Qy 1 CCGGGAGAGCCATAGTGGTCTGCG 24
| | | | | | | | | | : | | : | : | | |
Db 13 CCGGGAGAGCCAUAGUGGUCUGCG 36

Search completed: May 7, 2006, 06:32:34
Job time : 127.92 secs

SCORE 1.3 BuildDate: 12/06/2005 .

SCORE Search Results Details for Application 10759

Result us-10-759-514-6.rge.

[Score Home Page](#) [Retrieve Application List](#) [SCORE System Overview](#) [SCORE FAQ](#) [Comments / Suggestions](#)

This page gives you Search Results detail for the Application 10759514 and Search Result us-10-759-514-start

GenCore version 5.1.8
Copyright (c) 1993 - 2006 Bioceleration Ltd.

OM nucleic - nucleic search, using sw model

Run on: May 7, 2006, 06:16:29 ; Search time 512.074 Seconds
(without alignments)
2442.141 Million cell updates/sec

Title: US-10-759-514-6
Perfect score: 22
Sequence: 1 ttggcaacagtggcatgcaccg 22

Scoring table: IDENTITY_NUC
Gapop 10.0 , Gapext 1.0

Searched: 5883141 seqs, 28421725653 residues

Total number of hits satisfying chosen parameters: 11766282

Minimum DB seq length: 0
Maximum DB seq length: 2000000000

Post-processing: Minimum Match 0%
Maximum Match 100%
Listing first 45 summaries

Database : GenEmbl:*
1: gb_ba:*
2: gb_in:*
3: gb_env:*
4: gb_om:*
5: gb_ov:*
6: gb_pat:*
7: gb_ph:*
8: gb_pr:*
9: gb_ro:*
10: gb_sts:*
11: gb_sy:*
12: gb_un:*
13: gb_vi:*
14: gb_htg:*
15: gb_pl:*

Pred. No. is the number of results predicted by chance to have a score greater than or equal to the score of the result being printed, and is derived by analysis of the total score distribution.

SUMMARIES

%
Result Query

No.	Score	Match	Length	DB	ID	Description
c	1	22	100.0	1337	13 AF059603	AF059603 Wheat ros
c	2	22	100.0	5557	6 CQ830721	CQ830721 Sequence
c	3	22	100.0	5557	6 CQ832100	CQ832100 Sequence
c	4	22	100.0	48502	7 LAMCG	J02459 Bacterioph
c	5	19.4	88.2	44139	2 AY190942	AY190942 Drosophil
c	6	18.8	85.5	30844	15 AC158186	AC158186 Selagin
	7	18.8	85.5	104771	9 AL603830	AL603830 Mouse DNA
	8	18.8	85.5	206431	14 AC161885	AC161885 Gallus ga
	9	18.8	85.5	244279	14 AC163712	AC163712 Gallus ga
	10	18.8	85.5	248883	14 AC098544	AC098544 Rattus no
	11	18.8	85.5	259215	14 AC118088	AC118088 Rattus no
	12	18.8	85.5	262976	14 AC120483	AC120483 Rattus no
c	13	18.8	85.5	264908	9 AC096627	AC096627 Mus muscu
	14	18.4	83.6	110000	1 BA000023_21	Continuation (22 o
	15	18.4	83.6	110000	1 BA000023_22	Continuation (23 o
	16	18.4	83.6	145993	8 AC098972	AC098972 Homo sapi
	17	18.4	83.6	167304	9 AL928678	AL928678 Mouse DNA
c	18	18.4	83.6	169111	14 CR954168	CR954168 Danio rer
	19	18.4	83.6	197839	9 AL845466	AL845466 Mouse DNA
	20	18.4	83.6	207127	14 AC069496	AC069496 Homo sapi
c	21	18.4	83.6	213005	8 AP005059	AP005059 Homo sapi
	22	18.4	83.6	220618	14 AC131892	AC131892 Atelerix
c	23	18.4	83.6	237739	14 AC134520	AC134520 Atelerix
	24	17.8	80.9	905	10 BV576244	BV576244 G591P6073
c	25	17.8	80.9	32931	15 AC158190	AC158190 Selagin
	26	17.8	80.9	37165	15 AC158184	AC158184 Selagin
	27	17.8	80.9	39857	8 AC002522	AC002522 Homo sapi
	28	17.8	80.9	44873	8 AC004461	AC004461 Homo sapi
	29	17.8	80.9	59030	5 BX324184	BX324184 Zebrafish
	30	17.8	80.9	97351	8 AC015853	AC015853 Homo sapi
	31	17.8	80.9	108400	8 HUMDGCRCEN	L77570 Homo sapien
c	32	17.8	80.9	110000	1 BA000019_08	Continuation (9 of
	33	17.8	80.9	110000	15 CR382131_32	Continuation (33 o
	34	17.8	80.9	114540	8 AC107426	AC107426 Homo sapi
	35	17.8	80.9	125630	14 AC090650	AC090650 Arabidops
c	36	17.8	80.9	149939	5 AL935281	AL935281 Zebrafish
c	37	17.8	80.9	153284	8 AC078925	AC078925 Homo sapi
c	38	17.8	80.9	157327	14 CR450752	CR450752 Danio rer
	39	17.8	80.9	157904	8 AC108486	AC108486 Homo sapi
	40	17.8	80.9	159868	8 AC122129	AC122129 Homo sapi
c	41	17.8	80.9	159876	8 HS253P07	AL354000 Homo from
c	42	17.8	80.9	159994	8 AC157498	AC157498 Pan trogl
c	43	17.8	80.9	160256	5 BX950187	BX950187 Zebrafish
	44	17.8	80.9	160583	14 AC154944	AC154944 Monodelph
	45	17.8	80.9	169665	14 AC144566	AC144566 Homo sapi

ALIGNMENTS

RESULT 1

AF059603/c

LOCUS AF059603 1337 bp mRNA linear VRL 30-JAN-2000

DEFINITION Wheat rosette stunt virus nucleocapsid protein (N) mRNA, partial cds.

ACCESSION AF059603

VERSION AF059603.1 GI:6815246

KEYWORDS .

SOURCE Wheat rosette stunt virus

ORGANISM Wheat rosette stunt virus

Viruses; ssRNA negative-strand viruses; Mononegavirales;

Rhabdoviridae; unclassified Rhabdoviridae.

REFERENCE 1 (bases 1 to 1337)
AUTHORS Gong, Z.X.
TITLE Direct Submission
JOURNAL Submitted (15-APR-1998) Virology Laboratory, Shanghai Institute of Biochemistry, Chinese Academy of Science, 320 Yue Yang Rd., Shanghai 200031, P.R. China

FEATURES Location/Qualifiers
source 1. .1337
/organism="Wheat rosette stunt virus"
/mol_type="mRNA"
/db_xref="taxon:75890"
gene <1. .1337
/gene="N"
CDS <1. .1209
/gene="N"
/codon_start=1
/product="nucleocapsid protein"
/protein_id="AAF28467.1"
/db_xref="GI:6815247"
/translation="TFTKGDAERWVSTTKKASDSAFWLEVEGNSMTAPNGSKPSLAVI
VHLSSSVEVIFLGLSSSFRRSLICFRNYFIVILYHAYDNLLMSNLLIILNKLFSWLHQI
KSRMMLAKSHRHPLDCLFVCQESYSSYTFCHRLLLRHVIIRNLLAYRPIVFLIYKVD
DTVLRCTGDSMILRHIVIVHRIYIIIIICINRVACSATVAHRRYGRARRHVVCNVR
LAGELPIVRVLNDFQLLPILHIFCMREFVPPPTDHLSLYATVPRLLCAGATLQTSFSV
HATVANDLPRNWLASYRFECKNSPPHIKSIRSLPISVVISIFIIIFMIPRKNLPSSLAR
IQYSFPIIGKSEKSIIGFRFASINSDCSCRKRCGRITFPYNFYERVSLSNHFTQVLPC
LQTIPVSNIK"

ORIGIN

Query Match 100.0%; Score 22; DB 13; Length 1337;
Best Local Similarity 100.0%; Pred. No. 4.5;
Matches 22; Conservative 0; Mismatches 0; Indels 0; Gaps 0;

Qy 1 TTGGCAACAGTGGCATGCACCG 22
|||||
Db 845 TTGGCAACAGTGGCATGCACCG 824

RESULT 2
CQ830721/c
LOCUS CQ830721 5557 bp DNA linear PAT 12-JUL-2004
DEFINITION Sequence 24 from Patent WO2004055215.
ACCESSION CQ830721
VERSION CQ830721.1 GI:50251007
KEYWORDS .
SOURCE synthetic construct
ORGANISM synthetic construct
other sequences; artificial sequences.

REFERENCE 1
AUTHORS Otte, A.P. and van Blokland, H.J.
TITLE A method for improving protein production
JOURNAL Patent: WO 2004055215-A 24 01-JUL-2004;
Chromagenics B.V. (NL)

FEATURES Location/Qualifiers
source 1. .5557
/organism="synthetic construct"
/mol_type="unassigned DNA"
/db_xref="taxon:32630"
/note="lambda fragment 22425-27972"

ORIGIN

Query Match 100.0%; Score 22; DB 6; Length 5557;
Best Local Similarity 100.0%; Pred. No. 4;
Matches 22; Conservative 0; Mismatches 0; Indels 0; Gaps 0;

Qy 1 TTGGCAACAGTGGCATGCACCG 22
 |||||
 Db 1968 TTGGCAACAGTGGCATGCACCG 1947

RESULT 3

CQ832100/c

LOCUS CQ832100 5557 bp DNA linear PAT 29-JUL-2004

DEFINITION Sequence 9 from Patent WO2004056986.

ACCESSION CQ832100

VERSION CQ832100.1 GI:50831814

KEYWORDS .

SOURCE Bacteriophage lambda

ORGANISM Bacteriophage lambda

Viruses; dsDNA viruses, no RNA stage; Caudovirales; Siphoviridae;
 Lambda-like viruses.

REFERENCE 1

AUTHORS Otte,A.P., Kwaks,T.H. and Sewalt,R.G.

TITLE Means and methods for producing a protein through chromatin openers
 that are capable of rendering chromatin more accessible to
 transcription factors

JOURNAL Patent: WO 2004056986-A 9 08-JUL-2004;
 Chromagenics B.V. (NL)

FEATURES Location/Qualifiers

source 1. .5557
 /organism="Bacteriophage lambda"
 /mol_type="unassigned DNA"
 /db_xref="taxon:10710"

ORIGIN .

Query Match 100.0%; Score 22; DB 6; Length 5557;
 Best Local Similarity 100.0%; Pred. No. 4;
 Matches 22; Conservative 0; Mismatches 0; Indels 0; Gaps 0;

Qy 1 TTGGCAACAGTGGCATGCACCG 22
 |||||
 Db 1968 TTGGCAACAGTGGCATGCACCG 1947

RESULT 4

LAMCG/c

LOCUS LAMCG 48502 bp DNA circular PHG 17-APR-2002

DEFINITION Bacteriophage lambda, complete genome.

ACCESSION J02459 M17233 M24325 V00636 X00906

VERSION J02459.1 GI:215104

KEYWORDS DNA-binding protein; circular; coat protein; complete genome;
 origin of replication; repressor; unidentified reading frame.

SOURCE Bacteriophage lambda

ORGANISM Bacteriophage lambda

Viruses; dsDNA viruses, no RNA stage; Caudovirales; Siphoviridae;
 Lambda-like viruses.

REFERENCE 2 (bases 1 to 12)

AUTHORS Wu,R. and Taylor,E.

TITLE Nucleotide sequence analysis of DNA. II. Complete nucleotide
 sequence of the cohesive ends of bacteriophage lambda DNA

JOURNAL J. Mol. Biol. 57 (3), 491-511 (1971)

PUBMED 4931680

REFERENCE 3 (bases 45493 to 45963)

AUTHORS Imada,M. and Tsugita,A.

TITLE Amino acid sequence of lambda phage endolysin

JOURNAL Nature New Biol. 233, 230-231 (1971)

REFERENCE 4 (sites)

AUTHORS Weigel,P.H., Englund,P.T., Murray,K. and Old,R.W.

TITLE The 3'-terminal nucleotide sequences of bacteriophage lambda DNA
 JOURNAL Proc. Natl. Acad. Sci. U.S.A. 70 (4), 1151-1155 (1973)
 PUBMED 4515613
 REFERENCE 5 (bases 38597 to 38672)
 AUTHORS Dahlberg, J.E. and Blattner, F.R.
 TITLE In vitro transcription products of lambda DNA: Nucleotide sequences and regulatory sites
 JOURNAL (in) Fox, C.F. and Robinson, W.S. (Eds.);
 VIRUS RESEARCH. PROCEEDINGS OF 1973 ICN-UCLA SYMPOSIUM: 533-544;
 Academic Press, New York (1973)
 REFERENCE 6 (bases 37945 to 38027)
 AUTHORS Maniatis, T., Ptashne, M., Backman, K., Kield, D., Flashman, S.,
 Jeffrey, A. and Maurer, R.
 TITLE Recognition sequences of repressor and polymerase in the operators of bacteriophage lambda
 JOURNAL Cell 5 (2), 109-113 (1975)
 PUBMED 1095210
 REFERENCE 7 (bases 35583 to 35600)
 AUTHORS Kleid, D.G., Agarwal, K.L. and Khorana, H.G.
 TITLE The nucleotide sequence in the promoter region of the gene N in bacteriophage lambda
 JOURNAL J. Biol. Chem. 250 (14), 5574-5582 (1975)
 PUBMED 167018
 REFERENCE 8 (bases 35434 to 35618)
 AUTHORS Dahlberg, J.E. and Blattner, F.R.
 TITLE Sequence of the promoter-operator proximal region of the major leftward RNA of bacteriophage lambda
 JOURNAL Nucleic Acids Res. 2 (9), 1441-1458 (1975)
 PUBMED 1178525
 REFERENCE 9 (bases 37945 to 38018)
 AUTHORS Maniatis, T., Jeffrey, A. and Kleid, D.G.
 TITLE Nucleotide sequence of the rightward operator of phage lambda
 JOURNAL Proc. Natl. Acad. Sci. U.S.A. 72 (3), 1184-1188 (1975)
 PUBMED 1055375
 REFERENCE 10 (bases 44588 to 44773)
 AUTHORS Sklar, J., Yot, P. and Weissman, S.M.
 TITLE Determination of genes, restriction sites, and DNA sequences surrounding the 6S RNA template of bacteriophage lambda
 JOURNAL Proc. Natl. Acad. Sci. U.S.A. 72 (5), 1817-1821 (1975)
 PUBMED 1098044
 REFERENCE 11 (bases 37905 to 37989)
 AUTHORS Walz, A., Pirrotta, V. and Ineichen, K.
 TITLE Lambda repressor regulates the switch between PR and Prm promoters
 JOURNAL Nature 262 (5570), 665-669 (1976)
 PUBMED 958438
 REFERENCE 12 (bases 37946 to 38039)
 AUTHORS Smith, G.R., Eisen, H., Reichardt, L. and Hedgepeth, J.
 TITLE Deletions of lambda phage locating a prm mutation within the rightward operator
 JOURNAL Proc. Natl. Acad. Sci. U.S.A. 73 (3), 712-716 (1976)
 PUBMED 1062780
 REFERENCE 13 (bases 35578 to 35667; 37903 to 38027)
 AUTHORS Ptashne, M., Backman, K., Humayun, M.Z., Jeffrey, A., Maurer, R., Meyer, B. and Sauer, R.T.
 TITLE Autoregulation and function of a repressor in bacteriophage lambda
 JOURNAL Science 194 (4261), 156-161 (1976)
 PUBMED 959843
 REFERENCE 14 (bases 35578 to 35667)
 AUTHORS Humayun, Z., Jeffrey, A. and Ptashne, M.
 TITLE Completed DNA sequences and organization of repressor-binding sites in the operators of phage lambda
 JOURNAL J. Mol. Biol. 112 (2), 265-277 (1977)
 PUBMED 875019
 REFERENCE 15 (bases 38610 to 38732)

AUTHORS Scherer,G., Hobom,G. and Kossel,H.
 TITLE DNA base sequence of the po promoter region of phage lamdba
 JOURNAL Nature 265 (5590), 117-121 (1977)
 PUBMED 834253
 REFERENCE 16 (bases 38041 to 38241)
 AUTHORS Roberts,T.M., Shimatake,H., Brady,C. and Rosenberg,M.
 TITLE Sequence of Cro gene of bacteriophage lambda
 JOURNAL Nature 270 (5634), 274-275 (1977)
 PUBMED 593347
 REFERENCE 17 (bases 27616 to 28935)
 AUTHORS Davies,R.W., Schreier,P.H. and Buchel,D.E.
 TITLE Nucleotide sequence of the attachment site of coliphage lambda
 JOURNAL Nature 270 (5639), 757-760 (1977)
 PUBMED 593399
 REFERENCE 18 (bases 37206 to 37263; 37914 to 37970)
 AUTHORS Humayun,Z.
 TITLE DNA sequence at the end of the cI gene in bacteriophage lambda
 JOURNAL Nucleic Acids Res. 4 (7), 2137-2143 (1977)
 PUBMED 909767
 REFERENCE 19 (bases 27617 to 27934)
 AUTHORS Landy,A. and Ross,W.
 TITLE Viral integration and excision: structure of the lambda att sites
 JOURNAL Science 197 (4309), 1147-1160 (1977)
 PUBMED 331474
 REFERENCE 20 (bases 39062 to 39170)
 AUTHORS Denniston-Thompson,K., Moore,D.D., Kruger,K.E., Furth,M.E. and Blattner,F.R.
 TITLE Physical structure of the replication origin of bacteriophage lambda
 JOURNAL Science 198 (4321), 1051-1056 (1977)
 PUBMED 929187
 REFERENCE 21 (bases 44467 to 44807)
 AUTHORS Sklar,J.L.
 TITLE Structure and function of two regions of DNA controlling the synthesis of prokaryotic RNAs
 JOURNAL Thesis (1977)
 REFERENCE 22 (sites)
 AUTHORS Adhya,S. and Gottesman,M.
 TITLE Control of transcription termination
 JOURNAL Annu. Rev. Biochem. 47, 967-996 (1978)
 PUBMED 354508
 REFERENCE 23 (bases 13 to 72; 48391 to 48502)
 AUTHORS Nichols,B.P. and Donelson,J.E.
 TITLE 178-Nucleotide sequence surrounding the cos site of bacteriophage lambda DNA
 JOURNAL J. Virol. 26 (2), 429-434 (1978)
 PUBMED 666898
 REFERENCE 24 (bases 37938 to 38016; 35589 to 35666)
 AUTHORS Flashman,S.M.
 TITLE Mutational analysis of the operators of bacteriophage lambda
 JOURNAL Mol. Gen. Genet. 166 (1), 61-73 (1978)
 PUBMED 368570
 REFERENCE 25 (bases 37990 to 38982)
 AUTHORS Schwarz,E., Scherer,G., Hobom,G. and Kossel,H.
 TITLE Nucleotide sequence of cro, cII and part of the O gene in phage lambda DNA
 JOURNAL Nature 272 (5652), 410-414 (1978)
 PUBMED 264238
 REFERENCE 26 (bases 38212 to 38362)
 AUTHORS Rosenberg,M., Court,D., Shimatake,H., Brady,C. and Wulff,D.L.
 TITLE The relationship between function and DNA sequence in an intercistronic regulatory region in phage lambda
 JOURNAL Nature 272 (5652), 414-423 (1978)
 PUBMED 634366

REFERENCE 27 (bases 37224 to 37940)
AUTHORS Sauer,R.T.
TITLE DNA sequence of the bacteriophage gama cI gene
JOURNAL Nature 276 (5685), 301-302 (1978)
PUBMED 714163
REFERENCE 28 (bases 38597 to 39688)
AUTHORS Scherer,G.
TITLE Nucleotide sequence of the O gene and of the origin of replication
in bacteriophage lambda DNA
JOURNAL Nucleic Acids Res. 5 (9), 3141-3156 (1978)
PUBMED 704348
REFERENCE 29 (bases 29711 to 29811; 31043 to 31058)
AUTHORS Davies,R.W., Schreier,P.H. and Buchel,D.E.
TITLE Determination of the endpoints of partial deletion mutants of the
attachment site of bacteriophage lambda by DNA sequencing
JOURNAL Nucleic Acids Res. 5 (9), 3209-3218 (1978)
PUBMED 704352
REFERENCE 30 (bases 21661 to 31129)
AUTHORS Hoess,R.H. and Landy,A.
TITLE Structure of the lambda att sites generated by int-dependent
deletions
JOURNAL Proc. Natl. Acad. Sci. U.S.A. 75 (11), 5437-5441 (1978)
PUBMED 364480
REFERENCE 31 (bases 38453 to 38500)
AUTHORS Sprague,K.U., Faulds,D.H. and Smith,G.R.
TITLE A single base-pair change creates a Chi recombinational hotspot in
bacteriophage lambda
JOURNAL Proc. Natl. Acad. Sci. U.S.A. 75 (12), 6182-6186 (1978)
PUBMED 282634
REFERENCE 32 (bases 27711 to 27826)
AUTHORS Ross,W., Landy,A., Kikuchi,Y. and Nash,H.
TITLE Interaction of int protein with specific sites on lambda att DNA
JOURNAL Cell 18 (2), 297-307 (1979)
PUBMED 159130
REFERENCE 33 (bases 38008 to 39328)
AUTHORS Moore,D.D., Denniston-Thompson,K., Kruger,K.E., Furth,M.E.,
Williams,B.G., Daniels,D.L. and Blattner,F.R.
TITLE Dissection and comparative anatomy of the origins of replication of
lambdoid phages
JOURNAL Cold Spring Harb. Symp. Quant. Biol. 43 Pt 1, 155-163 (1979)
PUBMED 157834
REFERENCE 34 (bases 38470 to 39189)
AUTHORS Hobom,G., Grosschedl,R., Lusky,M., Scherer,G., Schwarz,E. and
Kossel,H.
TITLE Functional analysis of the replicator structure of lambdoid
bacteriophage DNAs
JOURNAL Cold Spring Harb. Symp. Quant. Biol. 43 Pt 1, 165-178 (1979)
PUBMED 157835
REFERENCE 35 (bases 38453 to 38500)
AUTHORS Smith,G.R., Faulds,D.H. and Sprague,K.U.

Query Match 100.0%; Score 22; DB 7; Length 48502;
Best Local Similarity 100.0%; Pred. No. 3.2;
Matches 22; Conservative 0; Mismatches 0; Indels 0; Gaps 0;

Qy 1 TTGGCAACAGTGGCATGCACCG 22
|||||
Db 24388 TTGGCAACAGTGGCATGCACCG 24367

RESULT 5
AY190942/c
LOCUS AY190942 44139 bp DNA linear INV 26-DEC-2002
DEFINITION Drosophila pseudoobscura clone DPSF01_33_N22 (D1412) genomic

sequence.

ACCESSION AY190942

VERSION AY190942.1 GI:27374267

KEYWORDS .

SOURCE Drosophila pseudoobscura

ORGANISM Drosophila pseudoobscura
Eukaryota; Metazoa; Arthropoda; Hexapoda; Insecta; Pterygota;
Neoptera; Endopterygota; Diptera; Brachycera; Muscomorpha;
Ephydroidea; Drosophilidae; Drosophila.

REFERENCE 1 (bases 1 to 44139)

AUTHORS Bergman,C.M., Pfeiffer,B.D., Rincon-Limas,D.E., Hoskins,R.A.,
Gnirke,A., Mungall,C.J., Wang,A.M., Kronmiller,B., Pacleb,J.,
Park,S., Stapleton,M., Wan,K., George,R.A., de Jong,P.J., Botas,J.,
Rubin,G.M. and Celniker,S.E.

TITLE Assessing the impact of comparative genomic sequence data on the
functional annotation of the Drosophila genome

JOURNAL Genome Biol. 3 (12), research0086 (2002)

REMARK http://genomebiology.com/2002/3/12/research/0086

REFERENCE 2 (bases 1 to 44139)

AUTHORS Pfeiffer,B.D., Bergman,C.M., Gnirke,A., Hoskins,R.A., Moshrefi,A.,
Mungall,C.J., Pacleb,J., Wang,A.M., Park,S., Wan,K., George,R.A.,
Rubin,G.M. and Celniker,S.E.

TITLE Direct Submission

JOURNAL Submitted (02-DEC-2002) Berkeley Drosophila Genome Project,
Lawrence Berkeley National Laboratory, 1 Cyclotron Rd, Berkeley, CA
94720, USA

COMMENT Gene annotations are preliminary because they are based solely on
alignments of conceptually translated gene models to the
corresponding gene model in Drosophila melanogaster. As a
consequence we did not annotate UTRs. In choosing splice sites, we
maximized conservation of amino acids rather than adherence to
consensus splice site.

FEATURES Location/Qualifiers

source 1. .44139
/organism="Drosophila pseudoobscura"
/mol_type="genomic DNA"
/strain="Tucson 14011-0121.4"
/db_xref="taxon:7237"
/clone="DPSF01_33_N22 (D1412)"
/clone_lib="DPSF01; partial MboI in pFOS1"

gene <108. .>1293
/gene="CG12130"

mRNA join(<108. .554,1129. .>1293)
/gene="CG12130"
/product="CG12130-PA"

CDS join(<108. .554,1129. .1293)
/gene="CG12130"
/codon_start=1
/product="CG12130-PA"
/protein_id="AAO01024.1"
/db_xref="GI:27374268"
/translation="GGELVIVNGPNAELGFHPDHYNEVHGFVMSMRNRQLVSKFGPNN
LEFKNPHDVAVTADGNEIYVAELNPMRIHKFLHKS LAKPMSLSATKDAQSISGVPAEA
PATGADASQTRHHPSGKAVLVASLMLIFAGSTFALALLMARRRRKRGCLPFGARSRRH
AWEKAEGFKLGGLLDRDRNGFEKLDQQASDEEQETLASGAPYA"

gene complement(<1722. .>2408)
/gene="CG1418"

mRNA complement(join(<1722. .2013,2081. .2287,2347. .>2408))
/gene="CG1418"
/product="CG1418-PA"

CDS complement(join(1722. .2013,2081. .2287,2347. .2408))
/gene="CG1418"
/codon_start=1
/product="CG1418-PA"

34779. .36668,36755. .37561,37643. .>37693)
 /gene="CG30421"
 /product="CG30421-PA"
 CDS join(25091. .25663,32116. .32202,34146. .34364,34442. .34711,
 34779. .36668,36755. .37561,37643. .37693)
 /gene="CG30421"
 /codon_start=1
 /product="CG30421-PA"
 /protein_id="AAO01028.1"
 /db_xref="GI:27374272"
 /translation="MGT TKEQQNEGDAAAEAAVGATHTTPTQHQRKETEEEEKTLP
 PVVASTMATNAAKKIRRAFSPMRNPFWRSRKFKTAAAAAIIASDAESGAGKDNAGERR
 GSHTAISSCVSGGRGRSGSIISLSSYEAPGDAAATTTNTYANSNSNHTNNTSNSN
 KRARVLRRSSFKKFLNRIQAHL SATVNVGVSSSEQSSAAASRVNLIDIVPFYTG CWRR
 KEANKHRQTNGLSASSGGGGDRVPPIGGYQWPPDQTPGVMGLKNHGNTCFMNAVLQC
 LSHTDILAEYFVLDQYKADLKRRNKINSRKFGTKGELTEQLANVLKALWTCRNE SDHS
 TGFKAVVD RYGTQFRSSTQHDAQEFLEFWLLDKVHEDLNTASKRRYKSIKQNSYGRSDE
 IIAAETLANHIRCNNSFVQAVFQAQFRSSLTCPRCKQSNTFDPFHCISVQLPQLTQQ
 TIFVTVVMKQQPRQIKMGLRVPAGSPIVALREQLQTDGTGIAGSRMVLVDLNAEGFTR
 VFYDTQPVETLSSIESIYCI E VPEAQAKIGATTKVAPSPSAPEAAEKIASAPAPAAP
 QADLLLLVANVYRAKD GKMTRF GAPFSMKAPRDCSYQDLQKRMLREMSPLLKPEVFS
 YTTPLADMFRIRLQDPSADPDTYLEQVEHPLLTEMIDMALSVLASEAGPPHIKLLLEW
 NSPEDHFLARQQETDDEAVVEHESVARLSASKAKDTAALTLEQCLEHYTKAETLSAED
 AWRCPHCQQYLPVVKTLGLWSLPDILVVHFKRFRQHQS KGPQAAKLTMTVKFPLTAFD
 MSPHLAKGVHESATSSSLGILDGGGRGSNLWKRARSGDSRSSTLNSRCDPKDTRYDLY
 AVCYHQGDTLETGHYTAACKNPYDRQWYKFDDQRVSKVPEDDIEQDIINNEAYMLFYQ
 RRSVDAGECSGSSSNSGDHWVSRIAPAPASATSSSADSTSGKENDGAIAEVSSEVA
 ESGPAAEKALSKPVNIPLVNEEGKRMSRISVTYMNINLYVILNIEELDFADADADVSL
 SEDAKNEGIEDPAEQIQTAAAAAAAAPKAAAIEEVTHIFAMDEEQKEEPVEVTKV
 ISRSEILPESLPTAEVVRTNGHVSPSSSSASSESSLSSRSSPRSLTTSVAAASTLQNK
 LNGHMNTACSSMFNGNGNGNANATHPQLSSSLQLTRQVLSNHNWHGSRSSVSAVENAT
 QQQKQAAAAASLSLRHSFSTSSNYKLRCTETLSSLLRNSTNTCSKDTLLFIDQQSH
 HHTAGLIEDDDSYMGRSLWISPVTPHKLITVSPKN"
 gene <40664. .>42391
 /gene="CG9196"
 mRNA join(<40664. .40677,40956. .41243,41317. .41652,
 41749. .>42391)
 /gene="CG9196"
 /product="CG9196-PA"
 CDS join(40664. .40677,40956. .41243,41317. .41652,41749. .42391)
 /gene="CG9196"
 /codon_start=1
 /product="CG9196-PA"
 /protein_id="AAO01029.1"
 /db_xref="GI:27374273"
 /translation="MQRSHFLLLAHLGLGQSQQSPATDYGEEQPPEGYYAFVES PNAV
 PPRVRPPPYTFVNADCKDVAAGKKS AVSVHNICGDLNKGQIPKNPLGQNVLGEPYPFE
 LIRNQTLKFLSKTLPVLKADDTLPKVTQIIRDEPVEQLDNNNIDRPYPGSGSTRIRRS
 VPDGIQSNEYSYFDPALDEEDGHHQQQQQQREQQQKRAAEERKPRKFCDGGGVFCTLY
 RAIQGD TGSAAAATPTAERREEVGPIRYEGPPTPCPAKVEYATPVFAKNYQGAWRYVV
 QIPYEGYFTQTVEVTRCIQARCHYLDGGCLSSPRWVSLVAEIFYPNAEDTVPTSSTT
 TQAPSVQDFQAYQQYLQKRAGVATASDGTSSSTGPNAGAGVAPADAHCDGHDELGCFFQ
 VRLYYDWFLIPGSCKCWRPDYFAKYVRRKSVADL"
 gene <43749. .>44081
 /gene="CG3880"
 mRNA <43749. .>44081
 /gene="CG3880"
 /product="CG3880-PA"
 CDS 43749. .>44081
 /gene="CG3880"
 /codon_start=1
 /product="CG3880-PA"
 /protein_id="AAO01030.1"
 /db_xref="GI:27374274"

/translation="MLRQKTSTSMALNTSDALNGNSPSQGQSLDMDALPKQFVLMSK
KLFDILDDKQSGYVRYVDIEKGWQDDGSKGLPRGVLDLRRVTPSCGLLSFERFCAGL
KLCLLRNQQ"

ORIGIN

Query Match 88.2%; Score 19.4; DB 2; Length 44139;
Best Local Similarity 95.2%; Pred. No. 73;
Matches 20; Conservative 0; Mismatches 1; Indels 0; Gaps 0;

Qy 2 TGGCAACAGTGGCATGCACCG 22
||||||| |||||
Db 5482 TGGCAACAGTGCCATGCACCG 5462

RESULT 6

AC158186/c

LOCUS AC158186 30844 bp DNA linear PLN 26-MAY-2005

DEFINITION Selaginella moellendorffii clone JGIASXY-5F17, complete sequence.

ACCESSION AC158186

VERSION AC158186.2 GI:66730719

KEYWORDS HTG.

SOURCE Selaginella moellendorffii

ORGANISM Selaginella moellendorffii
Eukaryota; Viridiplantae; Streptophyta; Embryophyta; Tracheophyta;
Lycopodiophyta; Isoetopsida; Selaginellales; Selaginellaceae;
Selaginella.

REFERENCE 1 (bases 1 to 30844)

AUTHORS DOE Joint Genome Institute and Stanford Human Genome Center.

TITLE Direct Submission

JOURNAL Unpublished

REFERENCE 2 (bases 1 to 30844)

AUTHORS DOE Joint Genome Institute.

TITLE Direct Submission

JOURNAL Submitted (09-MAR-2005) Production Genomics Facility, DOE Joint
Genome Institute, 2800 Mitchell Drive B100, Walnut Creek, CA
94598-1698, USA

REFERENCE 3 (bases 1 to 30844)

AUTHORS Stanford Human Genome Center.

CONSRM DOE Joint Genome Institute

TITLE Direct Submission

JOURNAL Submitted (26-MAY-2005) DOE Joint Genome Institute, 2800 Mitchell
Drive, Walnut Creek, CA 94598, USA

COMMENT On May 26, 2005 this sequence version replaced gi:60650325.

Draft Sequence Produced by DOE Joint Genome Institute
www.jgi.doe.gov

Finishing Completed at Stanford Human Genome Center
www-shgc.stanford.edu

Quality: Phrap Quality >=40 99.9% of Sequence;
Estimated Total Number of Errors is 0.

FEATURES Location/Qualifiers

source 1. .30844

/organism="Selaginella moellendorffii"

/mol_type="genomic DNA"

/db_xref="taxon:88036"

/clone="JGIASXY-5F17"

ORIGIN

Query Match 85.5%; Score 18.8; DB 15; Length 30844;
Best Local Similarity 90.9%; Pred. No. 1.5e+02;
Matches 20; Conservative 0; Mismatches 2; Indels 0; Gaps 0;

Qy 1 TTGGCAACAGTGGCATGCACCG 22
||||| |||||
Db 2812 TTGGCACCAGTGGCATGCTCCG 2791

RESULT 7
AL603830

LOCUS AL603830 104771 bp DNA linear ROD 09-FEB-2005

DEFINITION Mouse DNA sequence from clone RP23-467E19 on chromosome 11 Contains the Map2k3 gene for mitogen activated protein kinase kinase 3, the Gtlf3a gene for gene trap locus F3a, the Gtlf3b gene for gene trap locus F3b and two CpG islands, complete sequence.

ACCESSION AL603830

VERSION AL603830.7 GI:17017794

KEYWORDS HTG; CpG island; Gtlf3a; Gtlf3b; kinase; Map2K3.

SOURCE Mus musculus (house mouse)

ORGANISM Mus musculus
Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi; Mammalia; Eutheria; Euarchontoglires; Glires; Rodentia; Sciurognathi; Muridae; Murinae; Mus.

REFERENCE 1 (bases 1 to 104771)

AUTHORS Clark, S.

TITLE Direct Submission

JOURNAL Submitted (04-FEB-2005) Wellcome Trust Sanger Institute, Hinxton, Cambridgeshire, CB10 1SA, UK. E-mail enquiries: vega@sanger.ac.uk
Clone requests: clonerequest@sanger.ac.uk

COMMENT On Nov 20, 2001 this sequence version replaced gi:16944205.
The following abbreviations are used to associate primary accession numbers given in the feature table with their source databases:
Em:, EMBL; Sw:, SWISSPROT; Tr:, TREMBL; Wp:, WORMPEP; Information on the WORMPEP database can be found at
http://www.sanger.ac.uk/Projects/C_elegans/wormpep -----
Genome Center
Center: Wellcome Trust Sanger Institute
Center code: SC
Web site: <http://www.sanger.ac.uk>
Contact: vega@sanger.ac.uk

This sequence was finished as follows unless otherwise noted: all regions were either double-stranded or sequenced with an alternate chemistry or covered by high quality data (i.e., phred quality >= 30); an attempt was made to resolve all sequencing problems, such as compressions and repeats; all regions were covered by at least one subclone; and the assembly was confirmed by restriction digest, except on the rare occasion of the clone being a YAC.
Sequence from the Mouse Genome Sequencing Consortium whole genome shotgun may have been used to confirm this sequence. Sequence data from the whole genome shotgun alone has only been used where it has a phred quality of at least 30.
RP23-467E19 is from the RPCI-23 Mouse BAC Library constructed by the group of Pieter de Jong.
For further details see <http://www.chori.org/bacpac/home.htm>
VECTOR: pBACe3.6.

FEATURES

source	Location/Qualifiers 1. .104771 /organism="Mus musculus" /mol_type="genomic DNA" /db_xref="taxon:10090" /chromosome="11" /clone="RP23-467E19" /clone_lib="RPCI-23"
gene	complement(join(36672. .37719,38280. .38325,39423. .39562,42295. .42372,42712. .42839,44702. .44818,45877. .45996,46219. .46332,47130. .47178,47537. .47603,57212. .57429)) /gene="Map2k3" /locus_tag="RP23-467E19.1-002"
gene	complement(join(36672. .37719,38280. .38325,39423. .39562,

42295. .42372,42712. .42839,43921. .43972,44702. .44818,45877. .45996,46219. .46332,47130. .47178,47537. .47603,57212. .57429))

/gene="Map2k3"

/locus_tag="RP23-467E19.1-001"

mRNA complement(join(36672. .37719,38280. .38325,39423. .39562,42295. .42372,42712. .42839,43921. .43972,44702. .44818,45877. .45996,46219. .46332,47130. .47178,47537. .47603,57212. .57429))

/gene="Map2k3"

/locus_tag="RP23-467E19.1-002"

/product="mitogen activated protein kinase kinase 3"

/note="match: cDNAs: BC007467.1"

mRNA complement(join(36672. .37719,38280. .38325,39423. .39562,42295. .42372,42712. .42839,43921. .43972,44702. .44818,45877. .45996,46219. .46332,47130. .47178,47537. .47603,57212. .57429))

/gene="Map2k3"

/locus_tag="RP23-467E19.1-002"

/product="mitogen activated protein kinase kinase 3"

/note="match: ESTs: AA008333.1 AA433619.1 AV238394.1 AW106586.1 AW539252.1 BB858602.1 BF234702.1 BI555968.1 BY019272.1 BY226967.1 BY685384.1 BY710384.1 CA569861.1 CB947897.1 CB951524.1 W78467.1

match: cDNAs: AK008141.1 AK011002.1 AK093838.1 BC032478.1 D87116.1 U66839.1"

CDS complement(join(37636. .37719,38280. .38325,39423. .39562,42295. .42372,42712. .42839,43921. .43972,44702. .44818,45877. .45996,46219. .46332,47130. .47178,47537. .47603,57212. .57260))

/gene="Map2k3"

/locus_tag="RP23-467E19.1-002"

/standard_name="OTTMUSP00000006687"

/note="match: proteins: O09110 P46734"

/codon_start=1

/product="mitogen activated protein kinase kinase 3"

/protein_id="CAI25794.1"

/db_xref="GI:56237940"

/db_xref="InterPro:IPR000719"

/db_xref="InterPro:IPR002290"

/db_xref="InterPro:IPR008271"

/db_xref="InterPro:IPR011009"

/translation="MESPAASPPASLPQTKGKSKRKKDLRISCVSKPPVSNPTPPRNLDSRTFITIGDRNFEVEADDLVTISELGRGAYGVVEKVRHAQSGTIMAVKRIRATVNTQEQKRLMLDLINMRTVDCFYTVTFYGFALFREGDVWICMELMDTSLDKFYRKVLEKNMKIPEDILGEIAVSIVRALEHLHLSKLSVIHRDVKPSNVLINKEGHVKMCDGFIGSYLVDSVAKTMDAGCKPYMAPERINPELNQKGYNVKSDVWSLGITMIEMAILRFPYESWGTPFQQLKQVVVEEPSQLPADQFSPEFVDFTSQCLRKNPAERMSYLELMEHPFFTLHKTKKTDIAAFVKEILGEDS"

gene complement(join(37662. .37719,38280. .38821))

/gene="Map2k3"

/locus_tag="RP23-467E19.1-006"

mRNA complement(join(37662. .37719,38280. .38821))

/gene="Map2k3"

/locus_tag="RP23-467E19.1-006"

/product="mitogen activated protein kinase kinase 3"

/note="match: ESTs: BF467851.1"

gene complement(join(41892. .42372,42712. .42839,43921. .43972,44702. .44818,45877. .45996,46219. .46332,47130. .47178,47537. .47603,57212. .57429))

/gene="Map2k3"

/locus_tag="RP23-467E19.1-003"

mRNA complement(join(41892. .42372,42712. .42839,43921. .43972,44702. .44818,45877. .45996,46219. .46332,47130. .47178,47537. .47603,57212. .57429))

```

/gene="Map2k3"
/locus_tag="RP23-467E19.1-003"
/product="mitogen activated protein kinase kinase 3"
/note="match: ESTs: BB538636.2 BY724665.1
match: cDNAs: AK053378.1"
gene complement(join(42712. .42839,43921. .43972,44702. .44818,
45877. .45996,46219. .46332,47083. .47178,47537. .47603))
/locus_tag="RP23-467E19.1-004"
mRNA complement(join(42712. .42839,43921. .43972,44702. .44818,
45877. .45996,46219. .46332,47083. .47178,47537. .47603))
/locus_tag="RP23-467E19.1-004"
/product="mitogen activated protein kinase kinase 3"
/note="match: ESTs: BI525531.1"
gene complement(join(47367. .47603,57212. .57429))
/locus_tag="RP23-467E19.1-005"
mRNA complement(join(47367. .47603,57212. .57429))
/locus_tag="RP23-467E19.1-005"
/product="mitogen activated protein kinase kinase 3"
/note="match: ESTs: BB857429.1"
gene join(57616. .57917,67578. .68566)
/locus_tag="RP23-467E19.2-001"
mRNA join(57616. .57917,67578. .68566)
/locus_tag="RP23-467E19.2-001"
/product="gene trap locus F3a"
/note="match: ESTs: AI466976.1 AV085862.1 AV092279.1
AV253865.2 AV279917.1 BF099931.1 BY357260.1 BY709130.1
BY709606.1 BY709640.1
match: cDNAs: AK009371.1 AK010055.1 AK010107.1 BC031537.1"
gene join(74733. .74800,82483. .82601,83362. .83529)
/locus_tag="RP23-467E19.3-002"
mRNA join(74733. .74800,82483. .82601,83362. .83529)
/locus_tag="RP23-467E19.3-002"
/product="gene trap locus F3b"
/note="match: ESTs: BY176117.1 BY178036.1 BY190148.1"
gene join(75671. .75854,82483. .82601,83362. .87237)
/locus_tag="RP23-467E19.3-001"
mRNA join(75671. .75854,82483. .82601,83362. .87237)
/locus_tag="RP23-467E19.3-001"
/product="gene trap locus F3b"
/note="match: ESTs: AI413349.1 AI425923.1 BE308889.1
BE996115.1 BU152631.1 BY177308.1 BY196792.1 BY198579.1
BY202343.1 BY203840.1 BY747733.1
match: cDNAs: AK004720.2 AK080453.1 BC034332.1 BC038816.1"
CDS join(75758. .75854,82483. .82601,83362. .83478)
/locus_tag="RP23-467E19.3-002"
/standard_name="OTTMUSP00000006007"
/note="match: proteins: Q8N6N6 Q9DBW3"
/codon_start=1
/product="gene trap locus F3b"
/protein_id="CAI25795.1"
/db_xref="GI:56237941"
/translation="MAHATPPSALEQGGPIRVEHDRQRRQFSVRLNGCHDRAVLLYEY
VGKRIVDLQHTEVPDAYRGRGI AKHLAKAALDFVVEEDLKAHLTCWYIQKYVKENPLP

```

QYLERLQP"
polyA_signal 86401. .86406
/gene="Gtlf3b"
/locus_tag="RP23-467E19.3-001"
polyA_site 86427
/gene="Gtlf3b"
/locus_tag="RP23-467E19.3-001"

ORIGIN

Query Match 85.5%; Score 18.8; DB 9; Length 104771;
Best Local Similarity 90.9%; Pred. No. 1.4e+02;
Matches 20; Conservative 0; Mismatches 2; Indels 0; Gaps 0;

Qy 1 TTGGCAACAGTGGCATGCACCG 22
||||||| |
Db 6473 TTGGCAACAGTGCCATGCACAG 6494

RESULT 8

AC161885
LOCUS AC161885 206431 bp DNA linear HTG 21-MAY-2005
DEFINITION Gallus gallus chromosome UNKNOWN clone CH261-15B19, WORKING DRAFT
SEQUENCE, 8 unordered pieces.
ACCESSION AC161885
VERSION AC161885.1 GI:66392501
KEYWORDS HTG; HTGS_PHASE1; HTGS_DRAFT; HTGS_FULLTOP.
SOURCE Gallus gallus (chicken)
ORGANISM Gallus gallus
Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi;
Archosauria; Aves; Neognathae; Galliformes; Phasianidae;
Phasianinae; Gallus.
REFERENCE 1 (bases 1 to 206431)
AUTHORS Wilson,R.K.
TITLE The sequence of Gallus gallus clone
JOURNAL Unpublished
REFERENCE 2 (bases 1 to 206431)
AUTHORS Wilson,R.K.
TITLE Direct Submission
JOURNAL Submitted (21-MAY-2005) Genetics, Genome Sequencing Center, 4444
Forest Park Parkway, St. Louis, MO 63108, USA

COMMENT

----- Genome Center -----
Center: Washington University Genome Sequencing Center
Center code: WUGSC
Web site:http://genome.wustl.edu
Contact: submissions@watson.wustl.edu
----- Project Information -----
Center project name: J_AA015B19

----- Summary Statistics -----
Sequencing vector: M13; 0%
Sequencing vector: plasmid; 100%
Chemistry: Dye-primer ET; 0% of reads
Chemistry: Dye-terminator Big Dye; 100% of reads
Assembly program: Phrap; version 0.990319
Consensus quality: 203515 bases at least Q40
Consensus quality: 204721 bases at least Q30
Consensus quality: 205241 bases at least Q20

* NOTE: This is a 'working draft' sequence. It currently
* consists of 8 contigs. The true order of the pieces
* is not known and their order in this sequence record is
* arbitrary. Gaps between the contigs are represented as
* runs of N, but the exact sizes of the gaps are unknown.

* This record will be updated with the finished sequence
 * as soon as it is available and the accession number will
 * be preserved.

* 1 3949: contig of 3949 bp in length
 * 3950 4049: gap of unknown length
 * 4050 9460: contig of 5411 bp in length
 * 9461 9560: gap of unknown length
 * 9561 20832: contig of 11272 bp in length
 * 20833 20932: gap of unknown length
 * 20933 37073: contig of 16141 bp in length
 * 37074 37173: gap of unknown length
 * 37174 56455: contig of 19282 bp in length
 * 56456 56555: gap of unknown length
 * 56556 100723: contig of 44168 bp in length
 * 100724 100823: gap of unknown length
 * 100824 155191: contig of 54368 bp in length
 * 155192 155291: gap of unknown length
 * 155292 206431: contig of 51140 bp in length.

```
FEATURES
    source                Location/Qualifiers
                        1. .206431
                        /organism="Gallus gallus"
                        /mol_type="genomic DNA"
                        /db_xref="taxon:9031"
                        /chromosome="UNKNOWN"
                        /clone="CH261-15B19"
    misc_feature          1. .3949
                        /note="assembly_name:Contig2"
                        clone_end:T7
                        vector_side:right"
    gap                   3950. .4049
                        /estimated_length=unknown
    misc_feature          4050. .9460
                        /note="assembly_name:Contig3"
    gap                   9461. .9560
                        /estimated_length=unknown
    misc_feature          9561. .20832
                        /note="assembly_name:Contig4"
    gap                   20833. .20932
                        /estimated_length=unknown
    misc_feature          20933. .37073
                        /note="assembly_name:Contig5"
    gap                   37074. .37173
                        /estimated_length=unknown
    misc_feature          37174. .56455
                        /note="assembly_name:Contig6"
    gap                   56456. .56555
                        /estimated_length=unknown
    misc_feature          56556. .100723
                        /note="assembly_name:Contig7"
    gap                   100724. .100823
                        /estimated_length=unknown
    misc_feature          100824. .155191
                        /note="assembly_name:Contig8"
    gap                   155192. .155291
                        /estimated_length=unknown
    misc_feature          155292. .206431
                        /note="assembly_name:Contig9"
```

ORIGIN

Query Match 85.5%; Score 18.8; DB 14; Length 206431;
 Best Local Similarity 90.9%; Pred. No. 1.3e+02;
 Matches 20; Conservative 0; Mismatches 2; Indels 0; Gaps 0;

Qy 1 TTGGCAACAGTGGCATGCACCG 22

Db 26993 TTAGCAACAGTGGCATGCACAG 27014

RESULT 9

AC163712

LOCUS AC163712 244279 bp DNA linear HTG 13-JUN-2005

DEFINITION Gallus gallus chromosome UNKNOWN clone CH261-29M9, WORKING DRAFT
SEQUENCE, 32 unordered pieces.

ACCESSION AC163712

VERSION AC163712.1 GI:67515067

KEYWORDS HTG; HTGS_PHASE1; HTGS_DRAFT; HTGS_FULLLTOP.

SOURCE Gallus gallus (chicken)

ORGANISM Gallus gallus

Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi;
Archosauria; Aves; Neognathae; Galliformes; Phasianidae;
Phasianinae; Gallus.

REFERENCE 1 (bases 1 to 244279)

AUTHORS Wilson, R.K.

TITLE The sequence of Gallus gallus clone

JOURNAL Unpublished

REFERENCE 2 (bases 1 to 244279)

AUTHORS Wilson, R.K.

TITLE Direct Submission

JOURNAL Submitted (13-JUN-2005) Genetics, Genome Sequencing Center, 4444
Forest Park Parkway, St. Louis, MO 63108, USA

COMMENT

----- Genome Center -----

Center: Washington University Genome Sequencing Center

Center code: WUGSC

Web site: <http://genome.wustl.edu>

Contact: submissions@watson.wustl.edu

----- Project Information -----

Center project name: J_AA029M09

----- Summary Statistics -----

Sequencing vector: M13; 0%

Sequencing vector: plasmid; 100%

Chemistry: Dye-primer ET; 0% of reads

Chemistry: Dye-terminator Big Dye; 100% of reads

Assembly program: Phrap; version 0.990319

Consensus quality: 231430 bases at least Q40

Consensus quality: 235038 bases at least Q30

Consensus quality: 237275 bases at least Q20

* NOTE: This is a 'working draft' sequence. It currently
* consists of 32 contigs. The true order of the pieces
* is not known and their order in this sequence record is
* arbitrary. Gaps between the contigs are represented as
* runs of N, but the exact sizes of the gaps are unknown.
* This record will be updated with the finished sequence
* as soon as it is available and the accession number will
* be preserved.

* 1 1283: contig of 1283 bp in length
* 1284 1383: gap of unknown length
* 1384 2584: contig of 1201 bp in length
* 2585 2684: gap of unknown length
* 2685 3930: contig of 1246 bp in length
* 3931 4030: gap of unknown length
* 4031 5280: contig of 1250 bp in length
* 5281 5380: gap of unknown length
* 5381 7337: contig of 1957 bp in length
* 7338 7437: gap of unknown length
* 7438 9021: contig of 1584 bp in length

```

*      9022      9121: gap of unknown length
*      9122      11089: contig of 1968 bp in length
*     11090      11189: gap of unknown length
*     11190      13486: contig of 2297 bp in length
*     13487      13586: gap of unknown length
*     13587      15833: contig of 2247 bp in length
*     15834      15933: gap of unknown length
*     15934      17435: contig of 1502 bp in length
*     17436      17535: gap of unknown length
*     17536      19371: contig of 1836 bp in length
*     19372      19471: gap of unknown length
*     19472      22284: contig of 2813 bp in length
*     22285      22384: gap of unknown length
*     22385      25532: contig of 3148 bp in length
*     25533      25632: gap of unknown length
*     25633      30124: contig of 4492 bp in length
*     30125      30224: gap of unknown length
*     30225      33511: contig of 3287 bp in length
*     33512      33611: gap of unknown length
*     33612      36770: contig of 3159 bp in length
*     36771      36870: gap of unknown length
*     36871      42189: contig of 5319 bp in length
*     42190      42289: gap of unknown length
*     42290      46491: contig of 4202 bp in length
*     46492      46591: gap of unknown length
*     46592      50239: contig of 3648 bp in length
*     50240      50339: gap of unknown length
*     50340      54250: contig of 3911 bp in length
*     54251      54350: gap of unknown length
*     54351      58613: contig of 4263 bp in length
*     58614      58713: gap of unknown length
*     58714      67049: contig of 8336 bp in length
*     67050      67149: gap of unknown length
*     67150      74757: contig of 7608 bp in length
*     74758      74857: gap of unknown length
*     74858      82727: contig of 7870 bp in length
*     82728      82827: gap of unknown length
*     82828      93016: contig of 10189 bp in length
*     93017      93116: gap of unknown length
*     93117     105498: contig of 12382 bp in length
*    105499     105598: gap of unknown length
*    105599     119371: contig of 13773 bp in length
*    119372     119471: gap of unknown length
*    119472     132312: contig of 12841 bp in length
*    132313     132412: gap of unknown length
*    132413     154469: contig of 22057 bp in length
*    154470     154569: gap of unknown length
*    154570     175487: contig of 20918 bp in length
*    175488     175587: gap of unknown length
*    175588     197601: contig of 22014 bp in length
*    197602     197701: gap of unknown length
*    197702     244279: contig of 46578 bp in length.

```

```

FEATURES
  source      Location/Qualifiers
              1. .244279
              /organism="Gallus gallus"
              /mol_type="genomic DNA"
              /db_xref="taxon:9031"
              /chromosome="UNKNOWN"
              /clone="CH261-29M9"
  misc_feature 1. .1283
              /note="assembly_name:Contig16"
  gap          1284. .1383
              /estimated_length=unknown
  misc_feature 1384. .2584

```

```

        /note="assembly_name:Contig30"
gap      2585. .2684
        /estimated_length=unknown
misc_feature 2685. .3930
        /note="assembly_name:Contig31"
gap      3931. .4030
        /estimated_length=unknown
misc_feature 4031. .5280
        /note="assembly_name:Contig32"
gap      5281. .5380
        /estimated_length=unknown
misc_feature 5381. .7337
        /note="assembly_name:Contig34"
gap      7338. .7437
        /estimated_length=unknown
misc_feature 7438. .9021
        /note="assembly_name:Contig35"
gap      9022. .9121
        /estimated_length=unknown
misc_feature 9122. .11089
        /note="assembly_name:Contig36"
gap      11090. .11189
        /estimated_length=unknown
misc_feature 11190. .13486
        /note="assembly_name:Contig37"
gap      13487. .13586
        /estimated_length=unknown
misc_feature 13587. .15833
        /note="assembly_name:Contig38"
gap      15834. .15933
        /estimated_length=unknown
misc_feature 15934. .17435
        /note="assembly_name:Contig39"
gap      17436. .17535
        /estimated_length=unknown
misc_feature 17536. .19371
        /note="assembly_name:Contig40"
gap      19372. .19471
        /estimated_length=unknown
misc_feature 19472. .22284
        /note="assembly_name:Contig41"
gap      22285. .22384
        /estimated_length=unknown
misc_feature 22385. .25532
        /note="assembly_name:Contig42"
gap      25533. .25632
        /estimated_length=unknown
misc_feature 25633. .30124
        /note="assembly_name:Contig43"
gap      30125. .30224
        /estimated_length=unknown
misc_feature 30225. .33511
        /note="assembly_name:Contig44"
gap      33512. .33611
        /estimated_length=unknown
misc_feature 33612. .36770
        /note="assembly_name:Contig45"
gap      36771. .36870
        /estimated_length=unknown
misc_feature 36871. .42189
        /note="assembly_name:Contig46"
gap      42190. .42289
        /estimated_length=unknown
misc_feature 42290. .46491

```

```

        /note="assembly_name:Contig47"
gap      46492. .46591
        /estimated_length=unknown
misc_feature 46592. .50239
        /note="assembly_name:Contig48"
gap      50240. .50339
        /estimated_length=unknown
misc_feature 50340. .54250
        /note="assembly_name:Contig49"
gap      54251. .54350
        /estimated_length=unknown
misc_feature 54351. .58613
        /note="assembly_name:Contig50"
gap      58614. .58713

```

Query Match 85.5%; Score 18.8; DB 14; Length 244279;
 Best Local Similarity 90.9%; Pred. No. 1.3e+02;
 Matches 20; Conservative 0; Mismatches 2; Indels 0; Gaps 0;

```

Qy      1 TTGGCAACAGTGGCATGCACCG 22
        || |||||
Db      170397 TTAGCAACAGTGGCATGCACAG 170418

```

RESULT 10

AC098544

LOCUS AC098544 248883 bp DNA linear HTG 10-MAY-2003

DEFINITION Rattus norvegicus clone CH230-103A8, *** SEQUENCING IN PROGRESS
 ***, 5 unordered pieces.

ACCESSION AC098544

VERSION AC098544.6 GI:30521653

KEYWORDS HTG; HTGS_PHASE1; HTGS_DRAFT; HTGS_ENRICHED.

SOURCE Rattus norvegicus (Norway rat)

ORGANISM Rattus norvegicus

Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi;
 Mammalia; Eutheria; Euarchontoglires; Glires; Rodentia;
 Sciurognathi; Muroidea; Muridae; Murinae; Rattus.

REFERENCE 1 (bases 1 to 248883)

AUTHORS Muzny,D.Marie., Metzker,M.Lee., Abramzon,S., Adams,C., Alder,J.,
 Allen,C., Allen,H., Alsbrooks,S., Amin,A., Anguiano,D.,
 Anyalebechi,V., Aoyagi,A., Ayodeji,M., Baca,E., Baden,H.,
 Baldwin,D., Bandaranaike,D., Barber,M., Barnstead,M., Benahmed,F.,
 Biswalo,K., Blair,J., Blankenburg,K., Blyth,P., Brown,M.,
 Bryant,N., Buhay,C., Burch,P., Burrell,K., Calderon,E.,
 Cardenas,V., Carter,K., Cavazos,I., Ceasar,H., Center,A.,
 Chacko,J., Chavez,D., Chen,G., Chen,R., Chen,Y., Chen,Z., Chu,J.,
 Cleveland,C., Cockrell,R., Cox,C., Coyle,M., Cree,A., D'Souza,L.,
 Davila,M.L., Davis,C., Davy-Carroll,L., De Anda,C., Dederich,D.,
 Delgado,O., Denson,S., Deramo,C., Ding,Y., Dinh,H., Divya,K.,
 Draper,H., Dugan-Rocha,S., Dunn,A., Durbin,K., Duval,B., Eaves,K.,
 Egan,A., Escotto,M., Eugene,C., Evans,C.A., Falls,T., Fan,G.,
 Fernandez,S., Finley,M., Flagg,N., Forbes,L., Foster,M., Foster,P.,
 Fraser,C.M., Gabisi,A., Ganta,R., Garcia,A., Garner,T., Garza,M.,
 Gebregeorgis,E., Geer,K., Gill,R., Grady,M., Guerra,W., Guevara,W.,
 Gunaratne,P., Haaland,W., Hamil,C., Hamilton,C., Hamilton,K.,
 Harvey,Y., Havlak,P., Hawes,A., Henderson,N., Hernandez,J.,
 Hernandez,R., Hines,S., Hladun,S.L., Hodgson,A., Hogues,M.,
 Hollins,B., Howells,S., Hulyk,S., Hume,J., Idlebird,D., Jackson,A.,
 Jackson,L., Jacob,L., Jiang,H., Johnson,B., Johnson,R., Jolivet,A.,
 Karpathy,S., Kelly,S., Kelly,S., Khan,Z., King,L., Kovar,C.,
 Kowis,C., Kraft,C.L., Lebow,H., Levan,J., Lewis,L., Li,Z., Liu,J.,
 Liu,J., Liu,W., Liu,Y., London,P., Longacre,S., Lopez,J.,
 Lorensuhewa,L., Loulseged,H., Lozado,R.J., Lu,X., Ma,J.,
 Maheshwari,M., Mahindartne,M., Mahmoud,M., Malloy,K., Mangum,A.,

Mangum,B., Mapua,P., Martin,K., Martin,R., Martinez,E.,
Mawhiney,S., McLeod,M.P., McNeill,T.Z., Meenen,E.,
Milosavljevic,A., Miner,G., Minja,E., Montemayor,J., Moore,S.,
Morgan,M., Morris,K., Morris,S., Munidasa,M., Murphy,M., Nair,L.,
Nankervis,C., Neal,D., Newton,N., Nguyen,N., Norris,S.,
Nwaokelemeh,O., Okwuonu,G., Olarnpunsagoon,A., Pal,S., Parks,K.,
Pasternak,S., Paul,H., Perez,A., Perez,L., Pfannkoch,C.,
Plopper,F., Poindexter,A., Popovic,D., Primus,E., Pu,L.-L.,
Puazo,M., Quiroz,J., Rachlin,E., Reeves,K., Regier,M.A., Reigh,R.,
Reilly,B., Reilly,M., Ren,Y., Reuter,M., Richards,S., Riggs,F.,
Rives,C., Rodkey,T., Rojas,A., Rose,M., Rose,R., Ruiz,S.J.,
Sanders,W., Savery,G., Scherer,S., Scott,G., Shatsman,S., Shen,H.,
Shetty,J., Shvartsbeyn,A., Sisson,I., Sitter,C.D., Smajs,D.,
Sneed,A., Sodergren,E., Song,X.-Z., Sorelle,R., Sosa,J.,
Steimle,M., Strong,R., Sutton,A., Svatek,A., Tabor,P., Taylor,C.,
Taylor,T., Thomas,N., Thomas,S., Tingey,A., Trejos,Z., Usmani,K.,
Valas,R., Vera,V., Villasana,D., Waldron,L., Walker,B., Wang,J.,
Wang,Q., Wang,S., Warren,J., Warren,R., Wei,X., White,F.,
Williams,G., Willson,R., Wleczyk,R., Wooden,H., Worley,K.,
Wright,D., Wright,R., Wu,J., Yakub,S., Yen,J., Yoon,L., Yoon,V.,
Yu,F., Zhang,J., Zhou,J., Zhou,X., Zhao,S., Dunn,D., von
Niederhausern,A., Weiss,R., Smith,D.R., Holt,R.A., Smith,H.O.,
Weinstock,G. and Gibbs,R.A.

TITLE Direct Submission
JOURNAL Unpublished
REFERENCE 2 (bases 1 to 248883)
AUTHORS Worley,K.C.
TITLE Direct Submission
JOURNAL Submitted (24-OCT-2001) Human Genome Sequencing Center, Department
of Molecular and Human Genetics, Baylor College of Medicine, One
Baylor Plaza, Houston, TX 77030, USA
REFERENCE 3 (bases 1 to 248883)
AUTHORS Rat Genome Sequencing Consortium.
TITLE Direct Submission
JOURNAL Submitted (10-MAY-2003) Human Genome Sequencing Center, Department
of Molecular and Human Genetics, Baylor College of Medicine, One
Baylor Plaza, Houston, TX 77030, USA
COMMENT On May 10, 2003 this sequence version replaced gi:22855582.
The sequence in this assembly is a combination of BAC based reads
and whole genome shotgun sequencing reads assembled using Atlas
(<http://www.hgsc.bcm.tmc.edu/projects/rat/>). Each contig described
in the feature table below represents a scaffold in the Atlas
assembly (a 'contig-scaffold'). Within each contig-scaffold,
individual sequence contigs are ordered and oriented, and separated
by sized gaps filled with Ns to the estimated size. The sequence
may extend beyond the ends of the clone and there may be sequence
contigs within a contig-scaffold that consist entirely of whole
genome shotgun sequence reads. Both end sequences and whole genome
shotgun sequence only contigs will be indicated in the feature
table.
----- Genome Center
Center: Baylor College of Medicine
Center code: BCM
Web site: <http://www.hgsc.bcm.tmc.edu/>
Contact: hgsc-help@bcm.tmc.edu
----- Project Information
Center project name: GHJG
Center clone name: CH230-103A8
----- Summary Statistics
Assembly program: Atlas 3.0;
Consensus quality: 223541 bases at least Q40
Consensus quality: 226408 bases at least Q30
Consensus quality: 228553 bases at least Q20
Estimated insert size: 235340; sum-of-contigs estimation

Quality coverage: 6x in Q20 bases; sum-of-contigs estimation

* NOTE: Estimated insert size may differ from sequence length
* (see http://www.hgsc.bcm.tmc.edu/docs/Genbank_draft_data.html).
* NOTE: This is a 'working draft' sequence. It currently
* consists of 5 contigs. The true order of the pieces
* is not known and their order in this sequence record is
* arbitrary. Gaps between the contigs are represented as
* runs of N, but the exact sizes of the gaps are unknown.
* This record will be updated with the finished sequence
* as soon as it is available and the accession number will
* be preserved.
* 1 242485: contig of 242485 bp in length
* 242486 242585: gap of unknown length
* 242586 243738: contig of 1153 bp in length
* 243739 243838: gap of unknown length
* 243839 245069: contig of 1231 bp in length
* 245070 245169: gap of unknown length
* 245170 246652: contig of 1483 bp in length
* 246653 246752: gap of unknown length
* 246753 248883: contig of 2131 bp in length.

FEATURES Location/Qualifiers
source 1. .248883
/organism="Rattus norvegicus"
/mol_type="genomic DNA"
/db_xref="taxon:10116"
/clone="CH230-103A8"
misc_feature 1. .1709
/note="wgs_contig"
gap 242486. .242585
/estimated_length=unknown
gap 243739. .243838
/estimated_length=unknown
gap 245070. .245169
/estimated_length=unknown
gap 246653. .246752
/estimated_length=unknown

ORIGIN

Query Match 85.5%; Score 18.8; DB 14; Length 248883;
Best Local Similarity 90.9%; Pred. No. 1.3e+02;
Matches 20; Conservative 0; Mismatches 2; Indels 0; Gaps 0;

Qy 1 TTGGCAACAGTGGCATGCACCG 22
||||||| |
Db 70892 TTGGCAACAGTGCCATGCACAG 70913

RESULT 11

AC118088
LOCUS AC118088 259215 bp DNA linear HTG 19-NOV-2002
DEFINITION Rattus norvegicus clone CH230-30B23, WORKING DRAFT SEQUENCE.
ACCESSION AC118088
VERSION AC118088.5 GI:25087858
KEYWORDS HTG; HTGS_PHASE2; HTGS_DRAFT; HTGS_FULLTOP.
SOURCE Rattus norvegicus (Norway rat)
ORGANISM Rattus norvegicus
Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi;
Mammalia; Eutheria; Euarchontoglires; Glires; Rodentia;
Sciurognathi; Muroidea; Muridae; Murinae; Rattus.
REFERENCE 1 (bases 1 to 259215)
AUTHORS Muzny,D.Marie., Metzker,M.Lee., Abramzon,S., Adams,C., Alder,J.,
Allen,C., Allen,H., Alsbrooks,S., Amin,A., Anguiano,D.,
Anyalebechi,V., Aoyagi,A., Ayodeji,M., Baca,E., Baden,H.,

Baldwin,D., Bandaranaike,D., Barber,M., Barnstead,M., Benahmed,F.,
 Biswalo,K., Blair,J., Blankenburg,K., Blyth,P., Brown,M.,
 Bryant,N., Buhay,C., Burch,P., Burrell,K., Calderon,E.,
 Cardenas,V., Carter,K., Cavazos,I., Ceasar,H., Center,A.,
 Chacko,J., Chavez,D., Chen,G., Chen,R., Chen,Y., Chen,Z., Chu,J.,
 Cleveland,C., Cockrell,R., Cox,C., Coyle,M., Cree,A., D'Souza,L.,
 Davila,M.L., Davis,C., Davy-Carroll,L., De Anda,C., Dederich,D.,
 Delgado,O., Denson,S., Deramo,C., Ding,Y., Dinh,H., Divya,K.,
 Draper,H., Dugan-Rocha,S., Dunn,A., Durbin,K., Duval,B., Eaves,K.,
 Egan,A., Escotto,M., Eugene,C., Evans,C.A., Falls,T., Fan,G.,
 Fernandez,S., Finley,M., Flagg,N., Forbes,L., Foster,M., Foster,P.,
 Fraser,C.M., Gabisi,A., Ganta,R., Garcia,A., Garner,T., Garza,M.,
 Gebregeorgis,E., Geer,K., Gill,R., Grady,M., Guerra,W., Guevara,W.,
 Gunaratne,P., Haaland,W., Hamil,C., Hamilton,C., Hamilton,K.,
 Harvey,Y., Havlak,P., Hawes,A., Henderson,N., Hernandez,J.,
 Hernandez,R., Hines,S., Hladun,S.L., Hodgson,A., Hogues,M.,
 Hollins,B., Howells,S., Hulyk,S., Hume,J., Idlebird,D., Jackson,A.,
 Jackson,L., Jacob,L., Jiang,H., Johnson,B., Johnson,R., Jolivet,A.,
 Karpathy,S., Kelly,S., Kelly,S., Khan,Z., King,L., Kovar,C.,
 Kowis,C., Kraft,C.L., Lebow,H., Levan,J., Lewis,L., Li,Z., Liu,J.,
 Liu,J., Liu,W., Liu,Y., London,P., Longacre,S., Lopez,J.,
 Lorensuhewa,L., Loulseged,H., Lozado,R.J., Lu,X., Ma,J.,
 Maheshwari,M., Mahindartne,M., Mahmoud,M., Malloy,K., Mangum,A.,
 Mangum,B., Mapua,P., Martin,K., Martin,R., Martinez,E.,
 Mawhiney,S., McLeod,M.P., McNeill,T.Z., Meenen,E.,
 Milosavljevic,A., Miner,G., Minja,E., Montemayor,J., Moore,S.,
 Morgan,M., Morris,K., Morris,S., Munidasa,M., Murphy,M., Nair,L.,
 Nankervis,C., Neal,D., Newton,N., Nguyen,N., Norris,S.,
 Nwaokelemeh,O., Okwuonu,G., Olarnpunsagoon,A., Pal,S., Parks,K.,
 Pasternak,S., Paul,H., Perez,A., Perez,L., Pfannkoch,C.,
 Plopper,F., Poindexter,A., Popovic,D., Primus,E., Pu,L.-L.,
 Puazo,M., Quiroz,J., Rachlin,E., Reeves,K., Regier,M.A., Reigh,R.,
 Reilly,B., Reilly,M., Ren,Y., Reuter,M., Richards,S., Riggs,F.,
 Rives,C., Rodkey,T., Rojas,A., Rose,M., Rose,R., Ruiz,S.J.,
 Sanders,W., Savery,G., Scherer,S., Scott,G., Shatsman,S., Shen,H.,
 Shetty,J., Shvartsbeyn,A., Sisson,I., Sitter,C.D., Smajs,D.,
 Sneed,A., Sodergren,E., Song,X.-Z., Sorelle,R., Sosa,J.,
 Steimle,M., Strong,R., Sutton,A., Svatek,A., Tabor,P., Taylor,C.,
 Taylor,T., Thomas,N., Thomas,S., Tingey,A., Trejos,Z., Usmani,K.,
 Valas,R., Vera,V., Villasana,D., Waldron,L., Walker,B., Wang,J.,
 Wang,Q., Wang,S., Warren,J., Warren,R., Wei,X., White,F.,
 Williams,G., Willson,R., Wleczyk,R., Wooden,H., Worley,K.,
 Wright,D., Wright,R., Wu,J., Yakub,S., Yen,J., Yoon,L., Yoon,V.,
 Yu,F., Zhang,J., Zhou,J., Zhou,X., Zhao,S., Dunn,D., von
 Niederhausern,A., Weiss,R., Smith,D.R., Holt,R.A., Smith,H.O.,
 Weinstock,G. and Gibbs,R.A.

TITLE	Direct Submission
JOURNAL	Unpublished
REFERENCE	2 (bases 1 to 259215)
AUTHORS	Worley,K.C.
TITLE	Direct Submission
JOURNAL	Submitted (13-APR-2002) Human Genome Sequencing Center, Department of Molecular and Human Genetics, Baylor College of Medicine, One Baylor Plaza, Houston, TX 77030, USA
REFERENCE	3 (bases 1 to 259215)
AUTHORS	Rat Genome Sequencing Consortium.
TITLE	Direct Submission
JOURNAL	Submitted (19-NOV-2002) Human Genome Sequencing Center, Department of Molecular and Human Genetics, Baylor College of Medicine, One Baylor Plaza, Houston, TX 77030, USA
COMMENT	On Nov 19, 2002 this sequence version replaced gi:23265586. The sequence in this assembly is a combination of BAC based reads and whole genome shotgun sequencing reads assembled using Atlas (http://www.hgsc.bcm.tmc.edu/projects/rat/). Each contig described

in the feature table below represents a scaffold in the Atlas assembly (a 'contig-scaffold'). Within each contig-scaffold, individual sequence contigs are ordered and oriented, and separated by sized gaps filled with Ns to the estimated size. The sequence may extend beyond the ends of the clone and there may be sequence contigs within a contig-scaffold that consist entirely of whole genome shotgun sequence reads. Both end sequences and whole genome shotgun sequence only contigs will be indicated in the feature table.

----- Genome Center

Center: Baylor College of Medicine
Center code: BCM
Web site: <http://www.hgsc.bcm.tmc.edu/>
Contact: hgsc-help@bcm.tmc.edu

----- Project Information

Center project name: GUP0
Center clone name: CH230-30B23

----- Summary Statistics

Assembly program: Phrap; version 0.990329
Consensus quality: 238249 bases at least Q40
Consensus quality: 240765 bases at least Q30
Consensus quality: 242460 bases at least Q20
Estimated insert size: 245419; sum-of-contigs estimation
Quality coverage: 7x in Q20 bases; sum-of-contigs estimation

* NOTE: Estimated insert size may differ from sequence length
* (see http://www.hgsc.bcm.tmc.edu/docs/Genbank_draft_data.html).
* NOTE: This is a 'working draft' sequence. It currently
* consists of 1 contigs. Gaps between the contigs
* are represented as runs of N. The order of the pieces
* is believed to be correct as given, however the sizes
* of the gaps between them are based on estimates that have
* provided by the submittor.
* This sequence will be replaced
* by the finished sequence as soon as it is available and
* the accession number will be preserved.
* 1 259215: contig of 259215 bp in length.

FEATURES	Location/Qualifiers
source	1. .259215 /organism="Rattus norvegicus" /mol_type="genomic DNA" /db_xref="taxon:10116" /clone="CH230-30B23"
misc_feature	1. .1448 /note="wgs_end_extension clone_end:Sp6"
misc_feature	1577. .3317 /note="wgs_end_extension clone_end:Sp6"
misc_feature	3368. .4702 /note="wgs_end_extension clone_end:Sp6"
misc_feature	complement(5819. .6709) /note="clone_boundary clone_end:Sp6 site: end_sequence:BH290572"
misc_feature	complement(255901. .256586) /note="clone_boundary clone_end:T7 site: end_sequence:BH290550"

ORIGIN

Query Match 85.5%; Score 18.8; DB 14; Length 259215;
Best Local Similarity 90.9%; Pred. No. 1.3e+02;
Matches 20; Conservative 0; Mismatches 2; Indels 0; Gaps 0;

Qy 1 TTGGCAACAGTGGCATGCACCG 22
|||||||
Db 97487 TTGGCAACAGTGGCATGCACAG 97508

RESULT 12

AC120483

LOCUS AC120483 262976 bp DNA linear HTG 15-NOV-2002
DEFINITION Rattus norvegicus clone CH230-13K13, WORKING DRAFT SEQUENCE, 2
unordered pieces.

ACCESSION AC120483

VERSION AC120483.4 GI:25008120

KEYWORDS HTG; HTGS_PHASE1; HTGS_DRAFT; HTGS_FULLTOP.

SOURCE Rattus norvegicus (Norway rat)

ORGANISM Rattus norvegicus

Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi;
Mammalia; Eutheria; Euarchontoglires; Glires; Rodentia;
Sciurognathi; Muroidea; Muridae; Murinae; Rattus.

REFERENCE 1 (bases 1 to 262976)

AUTHORS

Muzny,D.Marie., Metzker,M.Lee., Abramzon,S., Adams,C., Alder,J.,
Allen,C., Allen,H., Alsbrooks,S., Amin,A., Anguiano,D.,
Anyalebechi,V., Aoyagi,A., Ayodeji,M., Baca,E., Baden,H.,
Baldwin,D., Bandaranaike,D., Barber,M., Barnstead,M., Benahmed,F.,
Biswal,K., Blair,J., Blankenburg,K., Blyth,P., Brown,M.,
Bryant,N., Buhay,C., Burch,P., Burrell,K., Calderon,E.,
Cardenas,V., Carter,K., Cavazos,I., Ceasar,H., Center,A.,
Chacko,J., Chavez,D., Chen,G., Chen,R., Chen,Y., Chen,Z., Chu,J.,
Cleveland,C., Cockrell,R., Cox,C., Coyle,M., Cree,A., D'Souza,L.,
Davila,M.L., Davis,C., Davy-Carroll,L., De Anda,C., Dederich,D.,
Delgado,O., Denson,S., Deramo,C., Ding,Y., Dinh,H., Divya,K.,
Draper,H., Dugan-Rocha,S., Dunn,A., Durbin,K., Duval,B., Eaves,K.,
Egan,A., Escotto,M., Eugene,C., Evans,C.A., Falls,T., Fan,G.,
Fernandez,S., Finley,M., Flagg,N., Forbes,L., Foster,M., Foster,P.,
Fraser,C.M., Gabisi,A., Ganta,R., Garcia,A., Garner,T., Garza,M.,
Gebregeorgis,E., Geer,K., Gill,R., Grady,M., Guerra,W., Guevara,W.,
Gunaratne,P., Haaland,W., Hamil,C., Hamilton,C., Hamilton,K.,
Harvey,Y., Havlak,P., Hawes,A., Henderson,N., Hernandez,J.,
Hernandez,R., Hines,S., Hladun,S.L., Hodgson,A., Hogue,M.,
Hollins,B., Howells,S., Hulyk,S., Hume,J., Idlebird,D., Jackson,A.,
Jackson,L., Jacob,L., Jiang,H., Johnson,B., Johnson,R., Jolivet,A.,
Karpathy,S., Kelly,S., Kelly,S., Khan,Z., King,L., Kovar,C.,
Kowis,C., Kraft,C.L., Lebow,H., Levan,J., Lewis,L., Li,Z., Liu,J.,
Liu,J., Liu,W., Liu,Y., London,P., Longacre,S., Lopez,J.,
Lorensuhewa,L., Loulseged,H., Lozado,R.J., Lu,X., Ma,J.,
Maheshwari,M., Mahindartne,M., Mahmoud,M., Malloy,K., Mangum,A.,
Mangum,B., Mapua,P., Martin,K., Martin,R., Martinez,E.,
Mawhiney,S., McLeod,M.P., McNeill,T.Z., Meenen,E.,
Milosavljevic,A., Miner,G., Minja,E., Montemayor,J., Moore,S.,
Morgan,M., Morris,K., Morris,S., Munidasa,M., Murphy,M., Nair,L.,
Nankervis,C., Neal,D., Newton,N., Nguyen,N., Norris,S.,
Nwaokelemeh,O., Okwuonu,G., Olarnpunsagoon,A., Pal,S., Parks,K.,
Pasternak,S., Paul,H., Perez,A., Perez,L., Pfannkoch,C.,
Plopper,F., Poindexter,A., Popovic,D., Primus,E., Pu,L.-L.,
Puazo,M., Quiroz,J., Rachlin,E., Reeves,K., Regier,M.A., Reigh,R.,
Reilly,B., Reilly,M., Ren,Y., Reuter,M., Richards,S., Riggs,F.,
Rives,C., Rodkey,T., Rojas,A., Rose,M., Rose,R., Ruiz,S.J.,
Sanders,W., Savery,G., Scherer,S., Scott,G., Shatsman,S., Shen,H.,
Shetty,J., Shvartsbeyn,A., Sisson,I., Sitter,C.D., Smajs,D.,
Sneed,A., Sodergren,E., Song,X.-Z., Sorelle,R., Sosa,J.,
Steimle,M., Strong,R., Sutton,A., Svatek,A., Tabor,P., Taylor,C.,

Taylor,T., Thomas,N., Thomas,S., Tingey,A., Trejos,Z., Usmani,K.,
 Valas,R., Vera,V., Villasana,D., Waldron,L., Walker,B., Wang,J.,
 Wang,Q., Wang,S., Warren,J., Warren,R., Wei,X., White,F.,
 Williams,G., Willson,R., Wleczyk,R., Wooden,H., Worley,K.,
 Wright,D., Wright,R., Wu,J., Yakub,S., Yen,J., Yoon,L., Yoon,V.,
 Yu,F., Zhang,J., Zhou,J., Zhou,X., Zhao,S., Dunn,D., von
 Niederhausern,A., Weiss,R., Smith,D.R., Holt,R.A., Smith,H.O.,
 Weinstock,G. and Gibbs,R.A.

TITLE Direct Submission
 JOURNAL Unpublished
 REFERENCE 2 (bases 1 to 262976)
 AUTHORS Worley,K.C.
 TITLE Direct Submission
 JOURNAL Submitted (07-MAY-2002) Human Genome Sequencing Center, Department
 of Molecular and Human Genetics, Baylor College of Medicine, One
 Baylor Plaza, Houston, TX 77030, USA
 REFERENCE 3 (bases 1 to 262976)
 AUTHORS Rat Genome Sequencing Consortium.
 TITLE Direct Submission
 JOURNAL Submitted (15-NOV-2002) Human Genome Sequencing Center, Department
 of Molecular and Human Genetics, Baylor College of Medicine, One
 Baylor Plaza, Houston, TX 77030, USA
 COMMENT On Nov 15, 2002 this sequence version replaced gi:23265426.
 The sequence in this assembly is a combination of BAC based reads
 and whole genome shotgun sequencing reads assembled using Atlas
 (<http://www.hgsc.bcm.tmc.edu/projects/rat/>). Each contig described
 in the feature table below represents a scaffold in the Atlas
 assembly (a 'contig-scaffold'). Within each contig-scaffold,
 individual sequence contigs are ordered and oriented, and separated
 by sized gaps filled with Ns to the estimated size. The sequence
 may extend beyond the ends of the clone and there may be sequence
 contigs within a contig-scaffold that consist entirely of whole
 genome shotgun sequence reads. Both end sequences and whole genome
 shotgun sequence only contigs will be indicated in the feature
 table.

----- Genome Center
 Center: Baylor College of Medicine
 Center code: BCM
 Web site: <http://www.hgsc.bcm.tmc.edu/>
 Contact: hgsc-help@bcm.tmc.edu
 ----- Project Information
 Center project name: GXNG
 Center clone name: CH230-13K13
 ----- Summary Statistics
 Assembly program: Phrap; version 0.990329
 Consensus quality: 242345 bases at least Q40
 Consensus quality: 245593 bases at least Q30
 Consensus quality: 247807 bases at least Q20
 Estimated insert size: 249832; sum-of-contigs estimation
 Quality coverage: 7x in Q20 bases; sum-of-contigs estimation

 * NOTE: Estimated insert size may differ from sequence length
 * (see http://www.hgsc.bcm.tmc.edu/docs/Genbank_draft_data.html).
 * NOTE: This is a 'working draft' sequence. It currently
 * consists of 2 contigs. The true order of the pieces
 * is not known and their order in this sequence record is
 * arbitrary. Gaps between the contigs are represented as
 * runs of N, but the exact sizes of the gaps are unknown.
 * This record will be updated with the finished sequence
 * as soon as it is available and the accession number will
 * be preserved.
 * 1 258816: contig of 258816 bp in length
 * 258817 258916: gap of unknown length
 * 258917 262976: contig of 4060 bp in length.

```

FEATURES                      Location/Qualifiers
    source                    1. .262976
                              /organism="Rattus norvegicus"
                              /mol_type="genomic DNA"
                              /db_xref="taxon:10116"
                              /clone="CH230-13K13"
    misc_feature              1. .1927
                              /note="wgs_end_extension
                              clone_end:Sp6"
    misc_feature              complement(5096. .5877)
                              /note="clone_boundary
                              clone_end:Sp6
                              site:
                              end_sequence:BH274933"
    misc_feature              137741. .139060
                              /note="wgs_contig"
    misc_feature              complement(256373. .257121)
                              /note="clone_boundary
                              clone_end:T7
                              site:
                              end_sequence:BH274931"
    misc_feature              257721. .258816
                              /note="wgs_end_extension
                              clone_end:T7"
    gap                        258817. .258916
                              /estimated_length=unknown

```

ORIGIN

```

Query Match      85.5%;   Score 18.8;   DB 14;   Length 262976;
Best Local Similarity  90.9%;   Pred. No. 1.3e+02;
Matches   20;   Conservative    0;   Mismatches    2;   Indels    0;   Gaps    0;

```

```

QY      1 TTGGCAACAGTGGCATGCACCG 22
        |||||
Db      45102 TTGGCAACAGTGCCATGCACAG 45123

```

RESULT 13

AC096627/c

LOCUS AC096627 264908 bp DNA linear ROD 03-MAY-2003

DEFINITION Mus musculus strain C57BL/6J clone rp23-77b23 map 11, complete sequence.

ACCESSION AC096627

VERSION AC096627.28 GI:30349025

KEYWORDS HTG.

SOURCE Mus musculus (house mouse)

ORGANISM Mus musculus

Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi; Mammalia; Eutheria; Euarchontoglires; Glires; Rodentia; Sciurognathi; Muroidea; Muridae; Murinae; Mus.

REFERENCE 1 (bases 1 to 264908)

AUTHORS Ni,Y., Song,L. and Roe,B.A.

TITLE Mus musculus BAC Clone rp23-77b23

JOURNAL Unpublished

REFERENCE 2 (bases 1 to 264908)

AUTHORS Ni,Y., Song,L. and Roe,B.A.

TITLE Direct Submission

JOURNAL Submitted (19-SEP-2001) Department Of Chemistry And Biochemistry, The University Of Oklahoma, 620 Parrington Oval, Room 208, Norman, OK 73019, USA

REFERENCE 3 (bases 1 to 264908)

AUTHORS Ni,Y., Song,L. and Roe,B.A.

TITLE Direct Submission

JOURNAL Submitted (02-MAY-2003) Department Of Chemistry And Biochemistry,

The University Of Oklahoma, 620 Parrington Oval, Room 208, Norman,
OK 73019, USA

REFERENCE 4 (bases 1 to 264908)

AUTHORS Ni,Y., Song,L. and Roe,B.A.

TITLE Direct Submission

JOURNAL Submitted (03-MAY-2003) Department Of Chemistry And Biochemistry,
The University Of Oklahoma, 620 Parrington Oval, Room 208, Norman,
OK 73019, USA

COMMENT On May 3, 2003 this sequence version replaced gi:29423919.
----- Genome Center
Center: Department Of Chemistry And Biochemistry
The University Of Oklahoma
Center code:UOKNOR
-----.

FEATURES Location/Qualifiers

source 1. .264908
/organism="Mus musculus"
/mol_type="genomic DNA"
/strain="C57BL/6J"
/db_xref="taxon:10090"
/map="11"
/clone="rp23-77b23"
/clone_lib="RPCI - 23 Female (C57BL/6J) Mouse BAC Library"

ORIGIN

Query Match 85.5%; Score 18.8; DB 9; Length 264908;
Best Local Similarity 90.9%; Pred. No. 1.3e+02;
Matches 20; Conservative 0; Mismatches 2; Indels 0; Gaps 0;

Qy 1 TTGGCAACAGTGGCATGCACCG 22
||||||| |
Db 172994 TTGGCAACAGTGCCATGCACAG 172973

RESULT 14
BA000023_21
WPCOMMENT

Sequence split into 27 fragments LOCUS BA000023 Accession BA000023

Fragment Name	Begin	End
BA000023_00	1	110000
BA000023_01	100001	210000
BA000023_02	200001	310000
BA000023_03	300001	410000
BA000023_04	400001	510000
BA000023_05	500001	610000
BA000023_06	600001	710000
BA000023_07	700001	810000
BA000023_08	800001	910000
BA000023_09	900001	1010000
BA000023_10	1000001	1110000
BA000023_11	1100001	1210000
BA000023_12	1200001	1310000
BA000023_13	1300001	1410000
BA000023_14	1400001	1510000
BA000023_15	1500001	1610000
BA000023_16	1600001	1710000
BA000023_17	1700001	1810000
BA000023_18	1800001	1910000
BA000023_19	1900001	2010000
BA000023_20	2000001	2110000
BA000023_21	2100001	2210000
BA000023_22	2200001	2310000
BA000023_23	2300001	2410000
BA000023_24	2400001	2510000

BA000023_25 2500001 2610000
BA000023_26 2600001 2694756
Continuation (22 of 27) of BA000023 from base 2100001 (BA000023 Sulfolobus tokodaii str. 7 DN

Query Match 83.6%; Score 18.4; DB 1; Length 110000;
Best Local Similarity 95.0%; Pred. No. 2.2e+02;
Matches 19; Conservative 0; Mismatches 1; Indels 0; Gaps 0;

Qy 2 TGGCAACAGTGGCATGCACC 21
|||||||
Db 100593 TGGCAACAGTGGGAATGCACC 100612

RESULT 15
BA000023_22
WPCOMMENT

Sequence split into 27 fragments LOCUS BA000023 Accession BA000023

Fragment Name	Begin	End
BA000023_00	1	110000
BA000023_01	100001	210000
BA000023_02	200001	310000
BA000023_03	300001	410000
BA000023_04	400001	510000
BA000023_05	500001	610000
BA000023_06	600001	710000
BA000023_07	700001	810000
BA000023_08	800001	910000
BA000023_09	900001	1010000
BA000023_10	1000001	1110000
BA000023_11	1100001	1210000
BA000023_12	1200001	1310000
BA000023_13	1300001	1410000
BA000023_14	1400001	1510000
BA000023_15	1500001	1610000
BA000023_16	1600001	1710000
BA000023_17	1700001	1810000
BA000023_18	1800001	1910000
BA000023_19	1900001	2010000
BA000023_20	2000001	2110000
BA000023_21	2100001	2210000
BA000023_22	2200001	2310000
BA000023_23	2300001	2410000
BA000023_24	2400001	2510000
BA000023_25	2500001	2610000
BA000023_26	2600001	2694756

Continuation (23 of 27) of BA000023 from base 2200001 (BA000023 Sulfolobus tokodaii str. 7 DN

Query Match 83.6%; Score 18.4; DB 1; Length 110000;
Best Local Similarity 95.0%; Pred. No. 2.2e+02;
Matches 19; Conservative 0; Mismatches 1; Indels 0; Gaps 0;

Qy 2 TGGCAACAGTGGCATGCACC 21
|||||||
Db 593 TGGCAACAGTGGGAATGCACC 612

Search completed: May 7, 2006, 07:20:01
Job time : 518.074 secs

SCORE 1.3 BuildDate: 12/06/2005